


- Salisbury, F.B. dan W.C. Ross. 1991. *Fisiologi Tumbuhan Jilid 2*. ITB. Bandung. Hal.32-35.
- Santi, L.P. dan D.H. Goenandi.2012. Efektivitas Dolomit Teraktivasi yang Diperkaya dengan Bakteri Pelarut Fosfat sebagai Pengganti Kiserit pada Bibit Kakao.*E-Journal Menara Perkebunan*. 8(1): 1-7.
- Sirenden, R.T., Suparno, dan Winerungan S.A.J. 2015. Hasil Tanaman Melon (*Cucumis melo* L.) Setelah Pemupukan Posfor dan Gandasil B pada Tanah Gambut Pedalaman.*Jurnal AGRI PEAT*. 16(1): 28-35.
- Suhartati, T. dan R. Kurniaty.2013. Inventarisasi Penyakit Daun pada Bibit di Stasiun Penelitian Nagrak.*Jurnal Perbenihan Tanaman Hutan*. 1(1): 51-59.
- Sumarni, N. dan R. Rosliani. 2001. Media Tumbuh dan Waktu Aplikasi Larutan Hara untuk Penanaman Cabai Secara Hidroponik. *J. Hort*. 11(4): 237-243.
- Sumenda, L., H.L. Rampe, dan F.R. Mantiri. 2011. Analisis Kandungan Klorofil Daun Mangga (*Mangifera indica* L.) pada Tingkat Perkembangan Daun yang Berbeda.*Jurnal Bioslogos*. 1(1): 20-24.
- Wulandari, E., B. Guritno, dan N. Aini. 2014. Pengaruh Kombinasi Jumlah Tanaman Per Polybag dan Komposisi Media Tanam terhadap Pertumbuhan dan Hasil Tanaman Mentimun (*Cucumis sativus* L.) Var. Venus. *Jurnal Protan*. 2(6): 464-473.
- Wijaya, K.A. 2008.*Nutrisi Tanaman sebagai Penentu Kualitas Hasil dan Resistensi Alami Tanaman*.Prestasi Pustaka Publisher. Jakarta. Hal.17-25.
- Yasmin, S., T. Wardiyati, dan Koesriharti.2014. Pengaruh Perbedaan Waktu Aplikasi dan Konesentrasi Giberelin (GA₃) terhadap Pertumbuhan dan Hasil Tanaman Cabai Besar.*Jurnal Produksi Tanaman*. 2(5):395-403.



**IMPACT OF DEHYDRATION ON PHYSIC-CHEMICAL PROPERTIES OF
YOGHURT AND LACTIC ACID BACTERIA ACTIVITY**

Ibrahim, A.I.^{1*}, Rifda, Naufalin², Tri, Yanto², Hidayah, Dwiyanti²

¹ Department of Food Science & Technology, Faculty of Agriculture Science, Omdurman Islamic University, P.O.BOX, www.oiu.edu.sd , Khartoum, Sudan

² Faculty of Agriculture, University of Jenderal Soedirman, UNSOED, Jl.dr. Soeparno Telp: (0281) 638791 Purwokerto 53123, Indonesia

^{*}) e-mail: ibrahim.sudann@gmail.com

ABSTRACT

Dehydration of food is the oldest operations used in food processing or food industry. Food dehydration is reducing moisture of food to low levels for increased shelf life. The yoghurt has rich to ash, also is contains nutrient. It is source of vitamins, phosphorus, potassium, calcium and protein, so considered suitable food for children and adults. It has contains lactic acid bacteria (LAB), that have function of health. Benefits of lactic acid bacteria represented in the lactose digestion, intestinal microflora modulation, cholesterol reduce cancer prevention and immune system stimulation. The yoghurt powder is produced by spray-drying at the optimum conditions, which can prolong a shelf life of it's to 90 days. Freeze-drying and air-drying can be used for dehydrating crispy dehydrated snacks. The freeze-drying lead to very aerated structure, softer final texture, and keeps original colour and flavour of raw materials. The air-drying lead to harder, crispier dried product and Mallard reactions. The freeze drying is suitable for yogurt dehydration because it lower effect on physico-chemical of product.

Keywords: *Yoghurt, Dehydration, Freeze-Drying Air-Drying Spry-Drying.*

1. INTRODUCTION

Milk is a source of nutritional compounds, water, fat, protein, lactose, minerals, enzymes, vitamins, and dissolved gases (Mourad et al, 2014). the composition of milk depend on several non-genetic factors(Contreras et al. 2015).

Cow's milk has highest percentage for protein, it has important role in human nutrition growth, and development. It has been the lowest fat content associated with digestive and heart problems. Cow milk is rich sources of minerals, vitamins and protein, such as calcium, vitamin D, riboflavin, and phosphorus , potassium, vitamin A, vitamin B12 and niacin (Ajai et al, 2012).

Goat's milk has contained higher amount of magnesium, phosphors and calcium than cow and human milks. It's important for prevention of diseases, and used for stimulation of immunity (Zenebe et al., 2014). Milk of goat has high nutritive value and therapeutic properties as a functional diet for human health (Abbas et al. 2014).

Soybean (*Glycine max*) is a one of plants leguminous. They are a rich source of good quality protein. Besides nutritional benefits soybeans provide several therapeutic benefits too. Soybean is contains all the major macronutrients required for good nutrition, as well vitamins, minerals, and fiber. in addition, Soybean protein has all the essential amino acids in the amounts needed for human health (Fan *et al*, 1995). Soy milk have been hypotensive in essential hypertensive patients (Rivas *et al*. 2002). Soy milk consumption has reduced blood pressure among diabetic patients with nephropathy (Miraghajani *et al*. 2013). Soybeans is contains almost 40% protein, thus soybeans

higher in protein than any other legumes and also many animal products. The composition of soybean seed are affects the soymilk and soy paneer quality and its yield (Poysa, 2002).

Lactic acid bacteria are a large group of bacteria widespread in nature and are beneficial in our digestive systems. They are the most important microorganisms used in food fermentation, contributing to the taste and texture and inhibiting food spoilage (Magdoub *et al*, 2015). The lactic acid bacteria starter cultures have several health, marketing, and technological advantages (Leroy, and Vuyst, 2004). Lactic acid bacteria in ripening cheese are added as starters and adjunct cultures or originate from the production and processing environments (Blaya *et al*, 2017). benefits of lactic acid bacteria represented in the lactose digestion, intestinal microflora modulation, cholesterol reduce, cancer prevention and immune system stimulation (Desobry-Banon *et al*, 1999).

Yoghurt is a fermented food product which is increasing its consumption by year, due to its benefit for human health (Agustini *et al*, 2017). Yogurts are one of most popular fermented dairy products widely consumed throughout the world (Loveday *et al*, 2013). The yoghurt has rich to ash, also is contains nutrient. It is source of vitamins, phosphorus, potassium, calcium and protein, so considered suitable food for children and adults (Igbabul *et al*, 2014). The total solids content, heat treatment, and incubation temperatures influence the textural properties of yogurts (Lee and Lucey, 2010). Cow and goat yoghurt has antioxidant activity and which is increasing during fermentation (Rahmawati and Suntornsuk, 2016). If yogurt of cow milk store at 5 °C, It can be safe until 14 days (Supavitpatana *et al*, 2010). The quality of yoghurt depend on three factors that are; the quality of raw milk, process conditions and starter culture (Paskov *et al*, 2010).Yoghurt can be mixed with variety of fruits, which are adding variety for the consumers (Kumar and Mishra, 2003). The enriched coconut yoghurt have been a good sensory in most quality attributes, compare with conventional yoghurt (Contreras *et al*. 2015).

The yoghurt powder produced by spray- drying at the optimum conditions. Which can prolong a shelf life of it's to 90 days by using packing material with lower moisture permeability and by selecting lower storage temperature and relative humidity (Balkir *et al*, 2011). The aim of this study was carried out to investigate the effect of dehydration on physic-chemical properties of yogurt and lactic acid bacteria activity, and increasing shelf-life of its.

2. YOGURT PREPARATION

The yogurt preparation according to method of (IDF,1987, (Kumar and Mishra, 2003)a, (Kumar and Mishra, 2003)b and (Zhao *et al.*, 2006). Preparation of raw milk with some modifications as follows: Cow's milk (milk fat 3%, total solids 12.3% and acidity 0.17%) was used for yoghurt production and 2% Skimmed milk powder was added to increase solids milk (Ebrahimi Sani and Islami, 2015). The Raw milk sample pasteurization separately at Pasteurization at 80-85 °C for 30 min or 90-95 °C for 5 min in stainless steel and cooled to 40°C. After cooling, starter (*Streptococcus*

thermophilus, *Lactobacillus bulgaricus* and *Lactobacillus acidophilus*) were inoculated at concentration of 3%, and incubated at 43 °C until the ph reached 4.5. (Miao *et al* , 2011).

3. YOGHURT MANUFACTURE

The yoghurt is manufactured according to international standards of yoghurt manufacture IDF standards (IDF,1987), and methods of (Kumar and Mishra, 2003)a, (Kumar and Mishra, 2003)b and (Zhao *et al.*, 2006). The milk is homogenized. And at 80-85 °C for 30 mints or 90-95 °C for 5 mints for pasteurization, and then cooled to 45°C. It is then inoculated with 2.5 or 3 present of a mixed lactic starter (2:1 *Streptococcus salivarius* ssp. *Thermophiles*, *Lactobacillus delbrueckii* ssp. *bulgaricus*), (*Streptococcus thermophilus*, *Lactobacillus bulgaricus* and *Lactobacillus acidophilus*).The inoculated milk is incubated to 45°C until a pH of 4.4 was attained in approximately 4 hours. The temperature increasing when incubation of yogurt, it act on increase viscosity and decreasing in values of ph (Wu and Zhang, 2016). The main processing steps in the manufacturing set and stirred yoghurt showed that in figure 1.




Figure 1. The main processing steps in the manufacturing set and stirred yoghurt
 Source: (Lee and Lucey, 2010)

4. YOGURT COMPOSITION

The yoghurt can be categorized into three major varieties, depend on the fat content of yogurt, it called regular yogurt, low-fat yogurt and non-fat yogurt. The regular yogurt is produced from

the full fat milk; it should be contain at least 3.25% of milk fat. low-fat yogurt and non-fat yogurt are producing from low fat milk or partially-skim milk, and skim milk sequences (Weerathilake *et al*, 2014). Yogurt is a dairy product obtained through the fermentation of milk, partly skimmed milk or skim milk by the lactic bacteria *Lactobacillus bulgaricus* and *Streptococcus thermophilus* with the lactic bacteria *Lactobacillus acidophilus*. The composition standards stipulate that yogurt must contain not less than 0.8% lactic acid, not less than 9.5% non-fat milk solids and not less than 3.0% protein

It may also contain some ingredients that come from milk (either whole or skim milk powder, or concentrated evaporated milk), fruits, fruit juices or extracts, jams, cereals or any other flavoring, sweeteners, a quantity not exceeding 2.0% of texturizing agents (stabilizers, gelling, thickening or emulsifying agents), citric acid, food coloring and, in the case of yogurt with added fruit, fruit juices or extracts or jams, a preservative not exceeding 50 ppm. By definition, a "fat-free" food must contain less than 0.5 g of fat per portion. (<http://www.milkingredients.ca/index-eng.php?id=197.9/3/2018>, 11:00). The composition of regular, low-fat, and non-fat yogurt showed that in table 1.

Table 1. The Composition of Regular, Low-Fat, and Non-Fat Yogurt

Parameter	Regular yogurt	Low-fat yogurt	Non-fat yogurt
Fat (%)	≥3.25	0.5 - 2.0	≤0.5
Solid Non Fat (%)	≥8.25	≥8.25	≥8.25
Titratable Acidity (%)	≥0.9	≥0.9	≥0.9
Ph	≤4.5	≤4.5	≤4.5

Sources: (Weerathilake *et al*, 2014).

5. ANTIMICROBIAL AND ANTIOXIDANT ACTIVITY

The Yogurt has contains LAB, that have function of health. Their growth depends on fermentation time, heat treatment and kind of sugar in milk (Hartati *et al*, 2012). Cow and goad yoghurt has antioxidant activity and which is increasing during fermentation (Rahmawati and Suntornsuk, 2016). Lactic acid bacteria can be producing antimicrobial compounds and it has contribution on human life (Khalid, 2011). Lactic acid bacteria have ability to degrade Ochratoxin A during the digestion in gastrointestinal and virto. Which can be used in silage industry, winemaking and starter culture during alcoholic fermentation (Luz *et al* , 2018). Lactic acid bacteria have ability on help digesting food in the intestine and fermenting the foods variety and turn them into tasty food products with increased shelf life (Teusink and Molenaar, 2017). Lactic acid bacteria have capacity to release peptides with antioxidant activity through proteolysis (Rizzello *et al*, 2017). the grape marc fermented by lactic acid bacteria and bifidobacteria strains can be used as pharmaceutical or food supplement (Campanella *et al*. 2017).

6. DEHYDRATION

Dehydration of food is the oldest operations used in food processing or food industry. Food dehydration is reducing moisture of food to low levels for increased shelf life. It is added a heat to the food by hot air, which also carries the moisture away from the food. Thus, it does not include removal of moisture from food by mechanical pressing or concentration of liquid foods (Jayas, 2016). Factors that affect stickiness have of great importance in the operation of spray dryers. Thus, stickiness of products during drying is effect on drying process, because product deposits in the drying chamber, product quality, product recovery, cleaning frequency (Refstrup *et al*, 2015).

Dehydration is the use hot air or heated surfaces for removes water from foods and reduce their water activity. This inhibits microbial growth and enzyme activity to extend their shelf-life (Fellows, 2017) a. The used heat for drying foods, or to concentrate liquids by boiling, removes water and therefore preserves the food by a reduction in water activity. However, the heat causes reduction in nutritional quality of product and effect on properties sensory. Also In freeze drying and freeze concentration is achieved preservation by reduction in water activity without heating the food with reduces the damage of heat (Fellows, 2017) b.

7. FREEZE DRYING

The Freeze drying is used in many applications since old years, most commonly in the food and pharmaceutical manufacture. Freeze drying depend on the removal of water from frozen product by a process called sublimation. The sublimation occurs when a frozen liquid goes directly to the gaseous state without passing through the liquid phase. The freeze drying process has been three stages: pre-freezing, primary drying, and secondary drying (Guide to Laboratory, 2010). The A typical phase diagram of freeze drying showed that in figure2.

8. YOGHURT DRYING

The prepared of dried yogurt samples were used method of (Carvalho *et al*, 2017). The formula for 100 grams of product is as follows: plain yogurts 42 g (Mercadona, Spain) were diverse with 25 g of mineral water. Separately, other 25g of water will be brought to a simmer and 4.4 g of is maltose, 2.2 g of malt dextrin (both from Cargill Iberica, Spain) and 0.2 g of sodium chloride was added until completely dissolved. 1.0 g of methylcellulose (Metolose MCE-400, Shin Etsu Chemical, Japan) and 0.2 g of xanthan gum (El Mazuelo, Spain) was then added and stirred. Immediately after, the mixture was added to the yogurt mix, mixed and put in a 1 L stainless steel foaming siphon, charged with one N₂O cartridge, shacked dynamically and directly put in a water/ice bath for 90 min. The siphon containing the blend was kept under cooling (4°C) for 4 days. After this resting period, the mixture (10 g) was whipped onto cupcake silicon moulds. Half the produced whipped yogurt foams were frozen down to -80°C for 24 hour, and was subsequently and immediately freeze-dried (FD) in a lab freeze-dryer (Lyoquest, Telstar, Spain) at 0.1 mbar for 22 hour. The rest of the foams were directly dried in a domestic dehydrator (Excalibur, Sacramento, USA), at 57°C for 24 hour. After

dehydration, keep the products into plastic sealed boxes with silica. The composition of the Commercial Yogurt Powder showed that in table 2, and number of lactic acid bacteria in yogurt powder (60°C) and fresh yogurt showed that in table 3.




Figure 2: A typical phase diagram
Source: (Guide to Laboratory, 2010).

Table 2. Composition of the Commercial Yogurt Powder

Component	Percentage (%)
Fat	1.2 - 2.0
Moisture	3.0 - 5.0
Protein	33.0-36.0
Ash	7.0 - 8.0
Lactose	50.0- 51.5

Source: (Krasaekoopt and Bhatia, 2012).

Table 3. The Number of Lactic Acid Bacteria in Yogurt Powder (60°C) and Fresh Yogurt

Yogurt type	Number of lactic acid bacteria (cfu g ⁻¹)
Fresh	9.8 x 10 ⁷
Powder	5.6 x 10 ⁷

Source: (Krasaekoopt and Bhatia, 2012).

9. EFFECT OF DEHYDRATION ON PRODUCTS

Power yoghurt produced by Spry-drying and Stored at 25 °C and 50% RH for 90 days their moisture and water activity values are increased. The titration acidity of samples is increased but PH value is decreased based on storage time. Also lactic acid bacteria count was decreased of storage

(Balkir *et al*, 2011). Freeze-drying and air-drying can be used for dehydrating crispy dehydrated snacks. Freeze-drying lead to very aerated structure, softer final texture, and keeps original colour and flavour of raw materials. The air-drying lead to harder and crispier dried product, Millard reactions (Carvalho *et al*, 2017). The Thermal stability studies that both processes of freeze-drying and spray-drying not effect in behaviour of whey protein(Vincenzetti *et al*. 2018). The drying conditions and the storage quality for dried milk is the very importance. Freeze-drying is good method than spray-drying and drum-drying for lower loss lysine from powder milk during storage (Aalaei *et al*, 2016). For obtain longer shelf life and good quality of product, the yoghurt powder can be stored at low temperatures and dry condition (Chutrtong 2015). Freeze drying and freeze concentration has capacity of reduce water activity of foods a with keep nutritional and sensory qualities (Khalid, 2011).

10. CONCLUSION

The yoghurt powder produced by spray- drying at the optimum conditions. Which can prolong a shelf life of it's to 90 days by using packing material with lower moisture permeability and by selecting lower storage temperature and relative humidity. Freeze-drying and air-drying can be used for dehydrating crispy dehydrated snacks. The freeze-drying lead to very aerated structure, softer final texture, and keeps original colour and flavour of raw materials. The air-drying lead to harder, crispier dried product and Mallard reactions. The freeze drying is suitable for yogurt dehydration because it lower effect on physico-chemical of product.

11. ACKNOWLEDGEMENT

The authors would like to thank of lecturers staff faculty of agriculture, Jenderal Soedirman University for their supports in completing the review.

12. REFERENCES

- Aalaei, Kataneh, Marilyn Rayner, and Ingegerd Sjöholm. 2016. "Storage Stability of Freeze-Dried, Spray-Dried and Drum-Dried Skim Milk Powders Evaluated by Available Lysine." *LWT - Food Science and Technology* 73:675–82. Retrieved (<http://dx.doi.org/10.1016/j.lwt.2016.07.011>).
- Abbas, H., F. A. Hassan, M. A. Abd El-Gawad, and A. Enab. 2014. "Physicochemical Characteristics of Goat's Milk." *Igarss 2014* 11(1):307–17.
- Agustini, T. W., D. Soetrisnanto, and W. F. Ma'ruf. 2017. "Study on Chemical, Physical, Microbiological and Sensory of Yoghurt Enriched by Spirulina Platensis." *International Food Research Journal* 24(1):367–71.
- Ajai, A.I; Ochigbo, S. S; Ndamitso, M. M and Olaoluwajuwon, J. 2012. "Proximate and Mineral Compositions of Different Raw Cow ' S Milks in Minna." 1(1):23–29.
- Balkir P., Sakin M.Koc B.Kaymak-Ertekin F. 2011. "Storage Stability of Spray-Dried Yoghurt Powder." *Milchwissenschaft* 66(2):159–62. Retrieved

- (<http://www.scopus.com/inward/record.url?eid=2-s2.0-79954615452&partnerID=40&md5=d903cda2eed27a870da65ad69b5d4795>).
- Blaya, J., Z. Barzideh, and G. LaPointe. 2017. "Interaction of Starter Cultures and Nonstarter Lactic Acid Bacteria in the Cheese Environment 1." *Journal of Dairy Science* 1–19. Retrieved (<http://linkinghub.elsevier.com/retrieve/pii/S0022030217311736>).
- Campanella, Daniela et al. 2017. "Exploitation of Grape Marc as Functional Substrate for Lactic Acid Bacteria and Bifidobacteria Growth and Enhanced Antioxidant Activity." *Food Microbiology* 65:25–35. Retrieved (<http://dx.doi.org/10.1016/j.fm.2017.01.019>).
- Carvalho, M. J., Perez-Palacios, T. and Ruiz-Carrascal, J. (2017) 'Physico-chemical and sensory characteristics of freeze-dried and air-dehydrated yogurt foam', *LWT - Food Science and Technology*. Elsevier Ltd, 80, pp. 328–334. doi: 10.1016/j.lwt.2017.02.039.
- Chutrtong, Jaruwan. 2015. "Survival of Probiotic Bacteria in Freeze - Dry Yogurt Starter Cultures Storage at 4 and 30 Degree Celsius." *Procedia - Social and Behavioral Sciences* 191:2219–25. Retrieved (<http://linkinghub.elsevier.com/retrieve/pii/S1877042815027743>).
- Contreras, Víctor Inocencio Pacheco, Gaspar Manuel Parra Bracamonte, Luis Arístides López Bustamante, Víctor Ricardo Moreno Medina, and Ana María Sifuentes Rincón. 2015. "Milk Composition and Its Relationship with Weaning Weight in Charolais Cattle." *Revista Brasileira de Zootecnia* 44(6):207–12.
- Desobry-Banon, S., N. Vétier, and J. Hardy. 1999. "Health Benefits of Yogurt Consumption. A Review." *International Journal of Food Properties* 2(1):1–12.
- Ebrahimi, A., Sani, A. M. and Islami, M. H. (2015) 'Evaluation of Rheological , Physicochemical , and Sensory Properties of Gundelia Tournefortii Yogurt', *Bulletin of Environment, Pharmacology and Life Sciences*, 3(3), pp. 213–231.
- Fan, M. Z., Sauer, W.C., de Lange, C.F.M. 1995. Amino acid digestibility in soyabean meal, extruded soyabean and full-fat canola for early-weaned pigs. *Anim. Feed Sci. Tech.* 52: 189-203.
- Fellows, P.J. 2017. 14 - Dehydration. *Food Processing Technology* (Fourth Edition). ISBN: 9780081019078. doi.org/10.1016/B978-0-08-100522-4.00014-6.
- Fellows, P.J. 2017. Freeze drying and freeze concentration. *Food Processing Technology*. ISBN: 9780081019078. DOI:10.1016/B978-0-08-100522-4.00023-7.
- Hartati, a I., Y. B. Pramono, and a M. Legowo. 2012. "Lactose And Reduction Sugar Concentrations, Ph And The Sourness Of Date Flavored Yogurt Drink As Probiotic Beverage." *Journal of Applied Food Technology* 1(1):1–3.
- Igbabul B., Shember J. and Amove J. 2014. "Physicochemical, Microbiological and Sensory Evaluation of Yoghurt Sold in Makurdi Metropolis." *African Journal of Food Science and Technology* 5(6):129–35. Retrieved (<http://dx.doi.org/10.14303/ajfst.2014.052>).
- International Dairy Federation. (1987) 'Skimmed milk whey and butter milk *determination of fat content. FIL-IDF Standard 22B, Bruxelles, Belgium.
- Jayas, Digvir, S. 2016. Food Dehydration. Reference Module in Food Science Principles. doi.org/10.1016/B978-0-08-100596-5.02913-9.
- Khalisanni Khalid. 2011. "An Overview of Lactic Acid Bacteria." *International Journal of Biosciences (IJB)* 1(3):1–13.
- Krasaekoopt, Wunwisa and Sumit Bhatia. 2012. "Production of Yogurt Powder Using Foam-Mat Drying." *AU Journal of Technology* 15(3):166–71.
- Kumar, P. and Mishra, H. N. .2003. 'Optimization of Mango Soy Fortified Yogurt Formulation Using Response Surface Methodology', *International Journal of Food Properties*, 6(3), pp. 499–517. doi: 10.1081/JFP-120021455.
- Kumar, Sanjeev, Birendra Kumar, Rajesh Kumar, and Suryamani Kumar. 2012. "Featur Eatures Nutritional F Eatur Es of Goat Milk Review Eview." *Indian Journal Dairy Science* 65(4)(July):266–73.
- Lee, W. J., and Lucey, J. A. 2010. ' Formation and Physical Properties of Yogurt. *Asian-Aust. J. Anim. Sci.* Vol. 23, No. 9 : 1127 – 1136.
- Leroy, Frédéric and Luc De Vuyst. 2004. "Lactic Acid Bacteria as Functional Starter Cultures for the Food Fermentation Industry." *Trends in Food Science and Technology* 15(2):67–78.

- Loveday, Simon M., Anwesha Sarkar, and Harjinder Singh. 2013. "Innovative Yoghurts: Novel Processing Technologies for Improving Acid Milk Gel Texture." *Trends in Food Science and Technology* 33(1):5–20.
- Luz, C., J. Ferrer, J. Mañes, and G. Meca. 2018. "Toxicity Reduction of Ochratoxin A by Lactic Acid Bacteria." *Food and Chemical Toxicology* 112(November 2017):60–66. Retrieved (<https://doi.org/10.1016/j.fct.2017.12.030>).
- Magdoub, Mohamed N. I. et al. 2015. "Probiotic Properties of Some Lactic Acid Bacteria Isolated from Egyptian Dairy Products." 4(12):758–66.
- Mourad, G., Bettache, G., and Samir M. 2014. 'Composition and nutritional value of raw milk', *Issues in Biological Sciences and Pharmaceutical Research* Vol.2 (10),pp .115-122. ISSN 2350-1588.
- Miao, Yu Zhi et al. 2011. "Extraction of Water-Soluble Polysaccharides (WSPS) from Chinese Truffle and Its Application in Frozen Yogurt." *Carbohydrate Polymers* 86(2):566–73. Retrieved (<http://dx.doi.org/10.1016/j.carbpol.2011.04.071>).
- Miraghajani, Maryam Sadat et al. 2013. "Soy Milk Consumption and Blood Pressure Among Type 2 Diabetic Patients With Nephropathy." *Journal of Renal Nutrition* 23(4):277–282.e1. Retrieved (<http://dx.doi.org/10.1053/j.jrn.2013.01.017>).
- Paskov, V., M. Karsheva, and I. Pentchev. 2010. "Effect of Starter Culture and Homogenization on the Rheological Properties of Yoghurts." *Journal of the University of Chemical Technology and Metallurgy* 1(January 2010):59–66.
- Poysa, V.2002. Stability of soybean seed composition and its effect on soymilk and tofu yield and quality. *Food Research International* 35: 337-345.
- Rahmawati, I. S. and Suntornasuk, W. (2016) 'Effects of Fermentation and Storage on Bioactive Activities in Milks and Yoghurts', *Procedia Chemistry*. Elsevier Ltd., 18(Mcls 2015), pp. 53–62. doi: 10.1016/j.proche.2016.01.010.
- Refstrup, E., Bonke, J., Ranadheera, CS. 2015. Plant and Equipment: Milk Dryers: Drying . Reference Module in Food Science-Principles, doi.org/10.1016/B978-0-08-100596-5.01012-X.
- Rivas, Miguel et al. 2002. "Soy Milk Lowers Blood Pressure in Men and Women with Mild to Moderate Essential Hypertension." *The Journal of Nutrition* 132(7):1900–1902.
- Rizzello, Carlo Giuseppe et al. 2017. "Improving the Antioxidant Properties of Quinoa Flour through Fermentation with Selected Autochthonous Lactic Acid Bacteria." *International Journal of Food Microbiology* 241:252–61. doi.org/10.1016/j.ijfoodmicro.2016.10.035).
- Supavitpatana, P., Wirjantoro, T. I. and Raviyan, P. (2010) 'Characteristics and shelf-life of Corn Milk Yogurt', *Chiang Mai University Journal of Natural Sciences*, 9(1), pp. 133–150.
- Teusink, Bas and Douwe Molenaar. 2017. "Systems Biology of Lactic Acid Bacteria: For Food and Thought." *Current Opinion in Systems Biology* 6:7–13. Retrieved (<http://linkinghub.elsevier.com/retrieve/pii/S2452310017301294>).
- Vincenzetti, Silvia et al. 2018. "Effects of Freeze-Drying and Spray-Drying on Donkey Milk Volatile Compounds and Whey Proteins Stability." *LWT - Food Science and Technology* 88:189–95. doi.org/10.1016/j.lwt.2017.10.019).
- Weerathilake, W.A., Rasika, D.M., Ruwanmali, J.K., and Munasinghe, M.A. 2014. The evolution, processing, varieties and health benefits of yogurt. *International Journal of Scientific and Research Publications*. ISSN 2250-3153.
- Wu, Hui, Liming Hu, and Guoping Zhang. 2016. "Effects of Electro-Osmosis on the Physical and Chemical Properties of Bentonite." *Journal of Materials in Civil Engineering* 28(8):6016010. Retrieved (<http://ascelibrary.org/doi/10.1061/%28ASCE%29MT.1943-5533.0001572>).
- Zhao, Qiang Zhong, Jin Shui Wang, Mou Ming Zhao, Yue Ming Jiang, and Cui Chun. 2006. "Effect of Casein Hydrolysates on Yogurt Fermentation and Texture Properties during Storage." *Food Technology and Biotechnology* 44(3):429–34.
- Zenebe, Tilahun, Nejash Ahmed, Tadele Kabeta, and Girma Kebede. 2014. "Review on Medicinal and Nutritional Values of Goat Milk." *Academic Journal of Nutrition* 3(3):30–39.

(<http://www.milkingredients.ca/index-eng.php?id=197.9/3/2018>, 11:00).



MENGENAL SENYAWA HCN PADA UBI KAYU

Kartika Noerwijati¹⁾ dan Rohmad Budiono²⁾

¹⁾Balai Penelitian Tanaman Aneka Kacang dan Umbi, Malang, Jawa Timur

²⁾Balai Pengkajian Teknologi Pertanian, Malang, Jawa Timur
email: tika_iletri@yahoo.com

ABSTRAK

Ubi kayu merupakan bahan makanan pokok ketiga setelah padi dan jagung, serta memiliki peran strategis dalam program diversifikasi pangan. Namun dalam pemanfaatannya sebagai bahan pangan, dibatasi oleh kandungan senyawa yang bersifat meracun jika dalam jumlah melebihi ambang batas. Senyawa tersebut adalah asam sianida atau Hydrogen sianide (HCN). Senyawa