

Gynura Procumbens Prevents Chemoresistance through Inhibition MDR1 Expression on MCF-7 Breast Cancer Cell Line and Sentizes the Cells to Doxorubicin

by Nunuk Aries Nurulita

Submission date: 06-Jul-2022 11:11AM (UTC+0700)

Submission ID: 1867166613

File name: 15998-30479-1-SM.pdf (284.41K)

Word count: 4202

Character count: 23115

3 ***Gynura procumbens* Prevents Chemoresistance through Inhibition MDR1 Expression on MCF-7 Breast Cancer Cell Line and Sensitizes the Cells to Doxorubicin**

Nunuk Aries Nurulita^{1,2*}, Edy Meiyanto³, Sugiyanto³, Eishou Matsuda⁴,
and Masashi Kawaichi⁴

¹ Faculty of Pharmacy Universitas Muhammadiyah Purwokerto, Purwokerto, Indonesia

² Graduate Program of Pharmaceutical Science, Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, Indonesia

³ Cancer Chemoprevention Research Centre, Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, Indonesia

⁴ Laboratorium of Gene Function of Animal, Nara Institute of Science and Technology, Nara, Japan

Abstract

The long-term exposure of doxorubicin (Dox) causes enhancement in MDR1 expression that leads to breast cancer cell resistance. This protein become a serious problem in cancer treatment and also well-known as negative prognostic factor in breast cancer malignancies. The new approach using natural chemopreventive substance was developed to inhibit this resistance progress. This study was aimed to investigate whether ethyl acetate fraction of *Gynura procumbens* (FEG) can prevent chemoresistance through suppressing the MDR1 protein expression. MCF-7 cell was used as chemoresistance cell model. The MCF-7 cells were maintained with 100 nM Dox-contained medium for five weeks. The chemoprevention effect of FEG was investigated by treated MCF-7/Dox with sub-toxic concentration of FEG. The cytotoxic properties of MCF-7 cells were determined by MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide) assay. Immunofluorescence and western blotting analysis was performed to detect the MDR1 expression. MCF-7/Dox cells need higher concentration for inhibiting cell growth, were compared with MCF-7, shown by IC₅₀ value. The MDR1 protein level elevated after Dox exposure in time dependent manner. The FEG treatment decreased MDR-1 protein level with dose dependent manner. FEG in combination with DOX potentiate the DOX effect on breast cancer cell growth inhibition. The FEG prevents the chemoresistance development in breast cancer cell line, MCF-7 induced by Dox through inhibiting MDR1 expression. The additional of FEG enhances Dox effect on cell death induction. Thus, FEG could be developed as co-chemotherapy agent for reverse multidrug resistance.

Keywords : *Gynura procumbens*, Chemoresistance, MDR1, MCF-7, MCF-7/Dox

Introduction

Drug resistance becomes a major obstacle to the successful of cancer chemotherapy. There are several mechanisms mediate the drug resistance mechanism including

drug inactivation, drug efflux, drug target mutation and the failure of apoptosis initiation (Davis *et al.*, 2003; Notarbartolo *et al.* 2005). Over expression of drug transporter such as P-gp plays a major role in causing multi drug resistance. P-gp is an adenosine triphosphatase (ATPase), a member of ATP-binding cassette (ABC) transporter encoded by MDR1 gene. In normal cells, P-gp localized at epithel cell and has a function to eliminate xenobiotic and metabolit endogen. However, in some human cancers P-gp overexpression has high correlation with the decreasing

¹⁴
*Corresponding author :

Nunuk Aries Nurulita,
Faculty of Pharmacy Universitas Muhammadiyah
Purwokerto, Jl. Raya Dukuwaluh, Kembaran,
Purwokerto, Jawa Tengah Indonesia
Telp. : +62-281-636751,
E-mail: nunuknurulita@yahoo.com

survival and become poor prognosis on cancer development (Leonessa and Clarke, 2003). There is positive correlation between overall *MDR1* expression and grade of breast cancer tumor (Surowiak *et al.*, 2005). This protein become a serious problem in cancer treatment and also well-known as negative prognostic factor in breast cancer malignancies.

P-gp substrates include substances in variable structure, chemical properties and drug mechanism. Several cancer chemotherapy drugs such as anthracyclines, taxanes, and epipodophylotoxins already well-known as the substrate of this protein. Long time exposure of these agents on cancer cells triggers the transcriptional activity of *MDR1* (30) and the increasing of P-gp (Germann, 1996; Nielsen *et al.*, 1996). The elevation of P-gp protein leads to chemotherapy failure due to cell resistance at several type of cancer. Doxorubicin which belongs to anthracyclin group could induce breast cancer multi drug resistance after long time exposure. DOX treatment increase P-gp expression level through Ras/Raf1 activation (Davis *et al.*, 2003). P-gp-mediated drug efflux decreases intracellular Dox uptake and decrease its efficacy. (48)

The development of P-gp inhibitors led to the third generation. Unfortunately, most of P-gp inhibitors that have been developed giving disappointing results. New approach using natural substances with moderate or low cytotoxic properties become a promising hope for reversing multi drug resistance due to *MDR1* over expression. Natural flavonoids have antitumor properties and others pharmacological activities related with its antitumor such as anti-inflammatory and antioxidant. Flavonoid, especially flavones and flavonol sh (36) modulation effect on cancer cell growth through cell cycle arrest and apoptosis induction (Limtrakul *et al.*, 2007). Quercetin and kaempferol down regulate multidrug resistance gene on human erythroleukemic K562/A cells. It causes decreasing of cells sensitivity to adriamycin

(Yanqiu *et al.*, 2011). Quercetin sensitizes cell resistance to daunorubicin through suppression on *MDR1* expression and P-gp activity at human pancreatic carcinoma (Borska *et al.*, 2010).

Gynura procumbens (28) is commonly used as herbal medicines in South-East Asia, especially Indonesia, Malaysia, and Thailand. (58) The extract and fraction of this plant have high anti-inflammatory and (37) antioxidant properties (Iskander *et al.*, 2002; Rosidah *et al.*, 2008). The ethyl acetate fraction of *G. procumbens* (FEG) may (53) contain quercetin and kaempferol. FEG sensitizes MCF-7 breast cancer cells to Doxorubicin treatment led to cell death. FEG modulates the microtubule integrity followed by cell cycle arrest and cell growth inhibition (Nurulita *et al.*, 2012). In this study, we subject FEG to doxorubicin-resistance cell. We use MCF-7 cells treated by 100 nM doxorubicin for five week as a model of cell resistance. We investigate the modulation on their sensitivity to doxorubicin, P-gp level, and cell growth.

Materials and Methods

Plant material

The leaves (26) *G. Procumbens* were obtained from Balai Penelitian dan Pengembangan Tanaman Obat dan Obat Tradisional (BP2TO2T) Indonesia. Dried powdered leaves were first extracted with ethanol 96%, and concentrated by evaporation under reduced pressure and the temperature was kept below 40°C. The extract was diluted in hot water and then fractionated with n-hexane. The aqueous fraction then was fractionated (21) in with ethyl acetate. The ethyl acetate fraction was concentrated by evaporation under reduced pressure and the temperature was kept below 40°C. The extract was stored at 4°C until used.

Cells and chemicals

MCF-7 (13) cells were obtained from Laboratorium of Gene Function in Animal, Nara Institute of Science and Technology (NAIST), Japan. MCF-7/DOX cells were

developed from MCF-7 cell that induced with 100 nM DOX for five weeks. The resistance development was checked by P-gp levels. The MCF-7 cells were routinely grown in DMEM (Nacalay Tesque) supplemented with 10% FBS (PAA Laboratories), 1% v/v Penicillin-streptomycin (Sigma), and 1 mM L-glutamine (Nacalay Tesque) at 37 °C in 5% CO₂. MCF-7/DOX cells were maintained in 25 nM DOX-contained medium. Doxorubicin (DOX) and 5-Fluorouracil (5-FU) (Ebewe) was purchased from PT. Ferron Par Pharmaceutical Cikarang, Indonesia, Dimethyl sulfoxide (DMSO) (Sigma, Aldrich, Germany), penicillin and streptomycin (Gibco), Tripsin (Sigma), 3-(4,5-dimethyliazol-2-il)-2,5-diphenyltetrazolium bromide (MTT) (Sigma), triton X-100, 4',6-diamidino-2-phenylindole (DAPI) (Invitrogen) are at analytical degree.

Antibodies

Primary antibodies that used in western blotting and immunostaining were anti P-Glycoprotein (JSB-1) (Genex, Inc.), anti α -Tubulin (Santa Cruz), and secondary antibodies were diluted in PBS containing 5% skim milk, 0,05% Tween (blocking buffer). For P-gp detection, the anti-P-gp and its secondary antibody were diluted in Can Get Signal (CGS) solution (Cosmo Bio).

Cells growth assay

Cells (10⁴ cells/well) were seeded 96-well plate (100 μ l/well). After 24 h of incubation, the medium was replaced with DOX, FEG and its combination-containing medium. The medium was discarded and replaced with MTT-containing medium (0.5 mg/ml) after 24 h incubation and incubated again for further 4 h at 37 °C, 5% CO₂. The reaction was stopped using 10% SDS in 0.1 N HCl solution and was incubated for overnight in light protected chamber, to dissolve formazan salt. The absorbance of each well was measured by ELISA reader at 595 nm. The ratio between treated and control cells absorbance referred to percentage (%) of cells viability.

Gel electrophoresis and immunoblotting

Cells were recovered, washed in PBS, and lysed for 30 min on ice using lysis buffer (20 mM Tris-HCl, pH 8, 0.5 mM EDTA, 1% NP40, 25 mM NaCl, and complete inhibitors of protease. Cells extract were centrifuged at 15.000 rpm for 20 min at 4°C to separate insoluble material. The protein concentration was determined using Bradford assay. Equal amount of each sample were mixed with SDS loading buffer, boiled for 3 min and subjected to SDS-PAGE at 120 V followed by electro blotting to Polyvinylidene fluoride (PVDF) membranes for 2 h at 100 V. Membranes were blocked using 5% skim milk in PBS at room temperature for 1 hour subsequently probed with the primary antibody of interest. Blots were exposed by Chemilumi-one Super (Nacalay Tesque).

Results

Long treatment of sub toxic concentration of DOX induced resistance cells development.


Breast cancer cells become less sensitive to chemotherapy agent during the cells resistance development. After five weeks induced by DOX, MCF-7 cells decreased its sensitivity to DOX treatment, as shown by IC₅₀ value (Table 1).

Table 1. IC₅₀ value of DOX and FEG on MCF-7 and MCF-7/DOX cells

Sample	Cell Line	IC ₅₀ value *
DOX	MCF-7	410 \pm 44 nM
	MCF-7/DOX	921 \pm 40 nM
FEG	MCF-7	259 \pm 29 μ g/ml
	MCF-7/DOX	161 \pm 40 μ g/ml

*average from two independent experiments

DOX resulted cell growth inhibition more than 2-fold higher on MCF-7/DOX sub line than that of MCF-7 cell line (Figure 1A). FEG treatment showed the opposite results. FEG caused cell growth inhibition stronger on MCF-7/DOX sub line compared with MCF-7 cell (Figure 1B).

5
A

B




Fig 2 1. MCF-7/DOX cells are less sensitive to DOX (A) but more sensitive to FEG (B). Cells (10^4 cells/well) were cultured at 96-well plate, then treated with indicated concentration of DOX or FEG for 24 h incubation. Cells Viability was determined with MTT assay. This result represents from 3 independent experiments. The data was statistically analyzed by two tailed-T-test analysis with $p < 0,05$.

DOX induced the elevation of P-gp level

DOX-induced MCF-7 cells showed the elevation of P-gp expression level in time dependent manner (Figure 2A). expression level of this protein was correlated with less sensitivity of MCF-7/DOX cells to DOX. FEG treatment decreased P-gp level in dose dependent pattern (Figure 2B). To make confirmation of this phenomenon, both immunoblotting and immunofluorescence were conducted. FEG decreased the green fluorescence intensity that refer P-gp protein decreasing (Figure 3). DOX induces breast cancer cell resistance due to increasing of

P-gp expression level. The P-gp protein has 170 kDa molecular size. This protein is hyperglycosylated as such appears as a diffuse band on Western blot results (Patel *et al.*, 2002). We also found the additional band below P-gp protein size (Figure 2). The lower molecules weight bands was detected using the JSB-1 antibody for P-gp represent degraded product of P-gp protein. JSB-1 antibody recognizes a cytoplasmic epitope of P-gp and does not cross-react with other protein such as MDR3 or muscle myosin, as does the C219 antibody (Leonessa and Clarke, 2003).




Figure 2. DOX induced P-gp expression on MCF-7 cells (A) and FEG prevents the elevation of this protein expression (B). Cells (10^6 cells/well) were seeded at 6-well plate, then treated with DOX, FEG or its combination for 24 h. Cell lysate then analyzed with Western blotting. Anti- α -tubulin was used for loading control. Dox1, Dox2, Dox3, and Dox4 refer to induction time for 2, 3, 4, and 5 weeks, respectively.

FEG reversed cell resistance and increased the sensitivity of MCF-7/DOX cells to DOX treatment

To investigate whether FEG could reverse the cell resistance development on MCF-7/DOX, MCF-7 cells were treated with DOX, FEG, or its combination and observed FEG reversal effect and its modulation on cell growth. Combination treatment of DOX+FEG suppressed cell growth of both of MCF-7 and MCF-7/DOX cells. The inhibition effect of combination treatment on MCF-7/DOX cells more significance compared to MCF-7 cells (Figure 4).

Discussion

P-glycoprotein (P-gp), a protein encoded by MDR-1 gene, has important function as ATP-dependent pump of drug efflux. P-gp has specific substrate, such as vinca alkaloids,

anthracycline, epipodophylotoxins, and rhodamine. The exposure of these substrate cancer cells could trigger transcriptional activity of MDR-1 (Germann, 1996; Lampidis *et al.*, 1989; Nielsen *et al.*, 1996).

Previous study have found the long time exposure of DOX induced cancer cells resistance through the increasing level of P-gp (Abolhoda *et al.*, 1999). In this study MCF-7 cells were used as resistance model of breast cancer. This cells were developed by long term sub toxic concentration of DOX induction. MCF-7 cells are cell line that express MDR1 and Breast cancer resistance protein (BCRP) that will be triggered its expression after several chemotherapy agent induction (Faneyte *et al.*, 2002). MDR-1 expression elevates at patient after long time chemotherapy agent administration and had strong correlation with cancer chemotherapy




Figure 3. DOX induced P-gp expression on pada MCF-7 cells and FEG prevents the elevation of this protein expression in concentration dependent manner. Cells ($5 \cdot 10^4$ cells/well) were seeded on cover slip at 24-well plate. After treated with or without DOX+FEG for 24 h, then stained with anti- α -Tubulin (green fluorescence) and counterstained with DAPI (blue fluorescence). Cells were observed under fluorescence microscope with 400x magnification.




Figure 4. FEG increased the sensitivity of MCF-7/DOX cells to DOX. Cells (10^4 cells/well) were cultured at 96-well plate, then treated with indicated concentration of DOX, FEG and its combination for 24 h incubation. Cells Viability was determined with MTT assay. This result represents from 10 independent experiments with similar results. The data was statistically analyzed by two tailed-T-test analysis. * $p = 0,1$ and ** $p = 0,028$.

failure (Leonessa and Clarke, 2003). This protein has identified as negative prognostic in cancer diseases development.

Sub toxic concentration of DOX for five weeks decreased MCF-7 cells sensitivity to this agent, shown by IC_{50} value become more than 2 fold higher compared with MCF-7 cells. DOX exposure to MCF-7 cells lead to increasing expression of P-gp in time dependent manner. MCF-7 cells after five weeks DOX induction (MCF-7/DOX) shown the significance increasing of this protein level compared with MCF-7 cells. The P-gp level has strong correlation with the low sensitivity of MCF-7/DOX to DOX exposure.

The attempts to overcome drug resistance mediated by *MDR-1* activation and elevation of P-gp level were developed the P-gp inhibitors. The agent that could suppress P-gp expression will increase uptake of chemotherapy agent and may influence its clinical efficacy. Flavonoids

could modulate the expression and activation of P-gp. There are several mechanisms mediate their effect: changing membrane cells permeability, suppressing P-gp expression, inhibition ATPase activity, and occupation substrate binding site. P-gp, then changes its conformation (Arora *et al.*, 2000; Boumendjel *et al.*, 2002; Drori *et al.*, 1995).

Natural compounds such as flavonoids and polyphenol have significant biological properties related to multidrug resistance development on cancer cells. Biochanin A and sylimarin inhibit drug efflux from the cells by P-gp pump, and produce synergistic effect when combined with DOX at MCF-7 and MDA435/LCC6. Both compound decreased IC_{50} value of daunomycin significantly at MCF-7/Adr cells (Zhang and Morris, 2003; Chung *et al.*, 2005). (-)-Epigallocatechin (EGC), (-)-epicatechin gallate (ECG), dan (-)-epigallocatechin gallate (EGCG) increase daunomycin accumulation at P-gp over-expressing cells, KB-C2 (Kitagawa *et al.*,

2004). Flavonoid and polyphenol have a promising properties to be developed as P-gp inhibitor to prevent cell resistance.

FEG treatment enhances DOX cytotoxicity significantly at MCF-7/DOX cells, higher than the effect on MCF-7 cells. The inhibition of cells growth exhibited synergistic effect of combination treatment DOX+FEG in dose dependent manner. FEG treatment decreases P-gp expression level at MCF-7/DOX cells linear to the given concentration. The ethyl acetate fraction of *G. procumbens* (FEG) was suggested have flavon and or flavonol with hydroxyl group modified to be metoksi at 3 and 4', and asetoksi at 7 position (Sugiyanto *et al.*, 2003). The ethyl acetate fraction of *G. procumbens* (FEG) may contain quercetin and kaempferol (Nurulita *et al.*, 2012). FEG may modulate *MDR-1* expression through the same mechanism with quercetin and kaempferol.

Quercetin inhibits the members of *MDR1* family, such as: P-gp, MRP, and BCRP, competitively (Cooray *et al.*, 2004; Scambia *et al.*, 1994; van Zanden *et al.*, 2005). Kaempferol and quercetin decrease P-gp expression and its activity at MDR KB-V1 cells (Limtrakul *et al.*, 2005). Quercetin and kaempferol prevent cell resistance development through down regulate multidrug resistance gene on human erythroleukemic K562/A cells. It reverses cells sensitivity to adriamycin (Yanqiu *et al.*, 2011). Quercetin suppresses *MDR1* expression and P-gp activity at human pancreatic carcinoma. The net influx of daunorubicin increases when given together with quercetin (Borska *et al.*, 2010). One study on structure activity relationship concludes flavonoid with -OH group at 5 and 7, with maximal have three total hydroxyl groups, and have strong inhibition on P-gp activity. Double bond at C2 and 3 as shown at flavon generates its planarity, lead to intercalate with hydrophobic amino acids residue of P-gp (Sheu *et al.*, 2010). Quercetin and kaempferol modulate both of *MDR-1* gene and P-gp protein level and also its activity that reverse multidrug resistance on many type of cancer cell lines.

This study found that FEG decrease P-gp expression level at MCF-7/DOX cells. This results were suggested to the decreasing P-gp expression that may affect on the increasing of DOX accumulation at cancer cells. In the next study, monitoring the net uptake of DOX during co-treatment of FEG should be considered. This effect may precede through several possibility, such as suppression on transcriptional activity of *MDR-1* gene, translation, or post translation.

As the conclusion, FEG prevents MCF-7 cells from developing resistance through down-regulating P-gp expression. FEG in combination with DOX potentiates the DOX effect on breast cancer cell growth inhibition. Thus, FEG could be developed as co-chemotherapy agent for reversing multidrug resistance.

Acknowledgement

This work was supported by Sandwich-like program 2010/2011, managing by Directorate General of Higher Education Republic of Indonesia, 2010

References

- Abolhoda, A., Wilson, A.E., Ross, H. Danenberg, P.V. Burt, M., and Scotto, K.W. 1999, Rapid activation of *MDR1* gene expression in human metastatic sarcoma after in vivo exposure to doxorubicin, *Clin. Cancer Res.*, **5**, 3352-3356
- Arora, A. Byrem, T.M., Nair, M.G., and Strasburg, G.M. 2000, Modulation of liposomal membrane fluidity by flavonoids and isoflavonoids, *Arch Biochem. Biophys.*, **373**, 102-109.
- Borska, S., Sopel, M., Chmielewska, M., Zabel, M., and Dziegiel, P. 2010, Quercetin as a potential modulator of P-glycoprotein expression and function in cells of human pancreatic carcinoma line resistant to daunorubicin, *Molecules*, **15**, 857-870.
- Boumendjel, A., Di Pietro, A., Dumontet, C., and Barron, D. 2002, Recent advances in the discovery of flavonoids and

- analogues with high affinity binding to P-glycoprotein responsible for cancer cell multidrug resistance, *Med. Res. Rev.*, **22**, 512-529.
- Chung, S.Y., Sung, M.K., Kim, N.H., Jang, J.O., Go, E.J., and Lee, H.J. 2005, Inhibition of P-glycoprotein by natural product in human breast cancer cells, *Arch. Pharm. Res.*, **28**(7), 823-828
- Cooray, H.C., Janvilisri, T., van Veen, H.W., Hladky, S.B., and Barrand, M.A. 2004, Interaction of the breast cancer resistance protein with plant polyphenols, *Biochem. Biophys. Res. Comm.*, **317**, 269-275
- Davis, J.M., Navolanic, P.M., Weinstein-Oppenheimer, C.R., Steelman, L.S., Wei H., Konopleva, M., Blagosklonny, M.V., and McCubrey, J.A. 2003, Raf-1 and Bcl-2 induce distinct and common pathways that contribute to breast cancer drug resistance, *Clin. Cancer Res.*, **9**, 1161-1170.
- Drori, S., Eytan, G.D., and Assaraf, Y.G. 1995, Potentiation of anticancer-drug Cytotoxicity by multidrug-resistance chemosensitizers involves alterations in membrane fluidity leading to increased membrane permeability, *Eur. J. Biochem.*, **228**, 1020-1029.
- Faneyte, I.F. Kristel, P.M.P. Maliepaard, M. Scheffer, G.L. Schepers, R.J. Schellens, J.H.M. and van de Vijver, 2002, Expression of the breast cancer resistance protein in breast cancer, *Clin. Cancer Res.*, **8**, 1068-1074.
- Germann, U.A. 1996, P-glycoprotein – mediator of multidrug resistance in tumor cells, *Eur. J. Cancer*. **32A**, 927-944
- Iskander, M.N., Song, Y., Coupar, I.M., and Jiratchariyakul, W. 2002, Antiinflammatory screening of medicinal plant *G. Procumbens*, *Plant Foods Hum. Nutr.*, **57**, 233-244.
- Kitagawa, S., Nabekura, T., and Kamiyama, S. 2004, Inhibition of P-glycoprotein function by tea catechins in KB-C2 cells. *J. Pharm. Pharmacol.*, **56**, 1001-1005.
- Leonessa, F. and Clarke, R. 2003, ATP binding cassette transporters and drug resistance in breast cancer, *Endocr. Relat. Cancer*, **10**, 43-73.
- Lampidis, T.J., Castello, C., del Giglio, A., Pressman, B.C., Viallet, P., Trevorrow, K.W., Valet, G.K., Tapiero, H., and Savaraj, N. 1989, Relevance of the chemical charge of rhodamine dyes to multiple drug resistance, *Biochem. Pharmacol.*, **38**, 4267-4271.
- Limtrakul, P., Khantamat, O., and Pintha, K. 2007, Inhibition of P-glycoprotein function and expression by kaempferol and quercetin. *J. Chemother.*, **17**, 86-95.
- Nielsen, D., Maare, C. and Skovsgaard, T. 1996, Cellular resistance to anthracyclins, *Gen. Pharmacol.*, **27**, 251-255.
- Notarbartolo, M., Poma, P., Perri, D., Dusonchet, L., Cervello, M., and Alessandro, N. 2005, Antitumor effects of curcumin, alone or in combination with cisplatin or doxorubicin, on human hepatic cancer cells. Analysis of their possible relationship to changes in NF- κ B activation levels and in IAP gene expression, *Cancer Lett.*, **224**, 53-65.
- Nurulita, N.A., Meiyanto, E., Sugiyanto, Matsuda, E., and Kawaichi, M., 2012, *Gynura procumbens* modulates the microtubules integrity and enhances distinct mechanism on doxorubicin and 5-fluorouracil-induced breast cancer cell death, *Orient. Pharm. Exp. Med.*, DOI 10.1007/s13596-012-0063-5.
- Nurulita, N.A., Meiyanto, E., and Sugiyanto, 2012, Selectivity of ethyl acetate fraction of *Gynura procumbens* on colon cancer and breast cancer, *Indo. J. Cancer Chemoprev.*, **3**(4)
- Pattel, V.A., Dunn, M.J., and Sorokin, A. 2002. Regulation of MDR-1 (P-glycoprotein) by cyclogenase 2. *J. Biol. Chem.*, **277**(41), 38915-38920.
- Rosidah, Yam, M.F., Sadikun, A. and Asmawi, M.Z. 2008, Antioxidant potential of *Gynura procumbens*, *Pharmaceutic. Biol.*, **46**, 616-625.

- Scambia, G., Ranelletti, F.O., Panici, P.B., De Vincenzo, R., Bonanno, G., Ferrandina, G., Piantelli, M., Bussa, S., Rumi, C., and Cianfriglia, M. 1994, Quercetin potentiates the effect of adriamycin in a multidrug-resistant MCF-7 human breast cancer cell line: P-glycoprotein as a possible target, *Cancer Chemother. Pharmacol.*, **34**, 459-464.
- Sheu, M.T., Liou, Y.B., Kao, Y.H., Lin, Y.K., and Ho, H.O. 2010, A quantitative structure-activity relationship for the modulation effects of flavonoids on P-glycoprotein-mediated transport, *Chem. Pharm. Bull.*, **58**(9), 1187-1194.
- Surowiak, P., Materna, V., Matkowski, R., Szczuraszek, K., Kornafel, J., Wojnar, A., Pudelko, M., Dietel, M., Denkert, C., Zabel, M., and Lage, H. 2005. Relationship between the expression of cyclooxygenase 2 and MDR1/P-glycoprotein in invasive breast cancers and their prognostic significance, *Breast Cancer Res.*, **7**, R862-R870.
- Sugiyanto, B., Sudarto, Meiyanto, E., Nugroho, A.E., and Jenie, U.A. 2003, The anticarcinogenic activity of plants compounds, *Indo. J. Pharm.*, **14**(4), 216-225.
- Van Zanden, J.J., Wortelboer, H. M., Bijlsma, S., Punt, A., Usta, M., Bladeren, P.J., Rietjens, I.M., and Cnubben, N.H. 2005, Quantitative structure activity relationship studies on the flavonoid mediated inhibition of multidrug resistance proteins 1 and 2. *Biochem. Pharmacol.*, **69**, 699-708, 2005.
- Yanqiu, H., Linjuan, C., Jin, W., Hongjun, H., Yongjin, S., Guobin, X., and Hanyun, R. 2011. The effect of quercetin and kaempferol on multidrug resistance and the expression of related genes in human erythroleukimia K562/A cells, *African J. Biotech*, **10**(62), 13399-13406.
- Zhang, S. and Morris, M.E. 2003, Effects of the flavonoids biochanin A, morin, phloretin, and silymarin on P-glycoprotein-mediated transport, *J. Pharmacol. Exp. Ther.*, **304**, 1258-1267.

Gynura Procumbens Prevents Chemoresistance through Inhibition MDR1 Expression on MCF-7 Breast Cancer Cell Line and Sentizes the Cells to Doxorubicin

ORIGINALITY REPORT

21 %
SIMILARITY INDEX

14 %
INTERNET SOURCES

17 %
PUBLICATIONS

3 %
STUDENT PAPERS

PRIMARY SOURCES

- 1** Sailendra N. Nichenametla, Todd G. Taruscio, Danny L. Barney, Jerry H. Exon. "A Review of the Effects and Mechanisms of Polyphenolics in Cancer", *Critical Reviews in Food Science and Nutrition*, 2006
Publication 1 %
- 2** brieflands.com
Internet Source 1 %
- 3** portalgaruda.ilkom.unsri.ac.id
Internet Source 1 %
- 4** etd.repository.ugm.ac.id
Internet Source 1 %
- 5** Sang, M.. "p73-dependent induction of 14-3-3@s increases the chemo-sensitivity of drug-resistant human breast cancers", *Biochemical and Biophysical Research Communications*, 20060818
Publication 1 %

6	Brand, W.. "Flavonoid-mediated inhibition of intestinal ABC transporters may affect the oral bioavailability of drugs, food-borne toxic compounds and bioactive ingredients", <i>Biomedicine & Pharmacotherapy</i> , 200611	1 %
Publication		
7	elifesciences.org	1 %
Internet Source		
8	David J. Villeneuve, Stacey L. Hembruff, Zachary Veitch, Melanie Cecchetto, William A. Dew, Amadeo M. Parissenti. "cDNA microarray analysis of isogenic paclitaxel- and doxorubicin-resistant breast tumor cell lines reveals distinct drug-specific genetic signatures of resistance", <i>Breast Cancer Research and Treatment</i> , 2005	1 %
Publication		
9	Submitted to Far Eastern University	1 %
Student Paper		
10	qmro.qmul.ac.uk	1 %
Internet Source		
11	www.mdpi.com	<1 %
Internet Source		
12	"Activity of inhibit the cell cycle and induct apoptosis in HeLa cancer cell with combination of Sabrang onion (Eleutherine	<1 %

palmifolia (L.) Merr) and Starfruit Mistletoe (Macrosolen cochinchinensis (Lour.) Tiegh)", Journal of Applied Pharmaceutical Science, 2018

Publication

13

Submitted to Program Pascasarjana Universitas Negeri Yogyakarta

Student Paper

<1 %

14

journal.uad.ac.id

Internet Source

<1 %

15

Jing Jin, Feng-Peng Wang, Huailing Wei, Gengtao Liu. "Reversal of multidrug resistance of cancer through inhibition of P-glycoprotein by 5-bromotetrandrine", Cancer Chemotherapy and Pharmacology, 2004

Publication

<1 %

16

lib.ugent.be

Internet Source

<1 %

17

Submitted to King's College

Student Paper

<1 %

18

Schutte, M.E.. "An in vitro and in silico study on the flavonoid-mediated modulation of the transport of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) through Caco-2 monolayers", Toxicology and Applied Pharmacology, 20061201

Publication

<1 %

19 Daniela Damiani. "P-glycoprotein, lung resistance-related protein and multidrug resistance-associated protein in de novo adult acute lymphoblastic leukaemia", *British Journal of Haematology*, 3/2002
Publication

20 Castro, A.F.. "Inhibition of drug transport by genistein in multidrug-resistant cells expressing P-glycoprotein", *Biochemical Pharmacology*, 19970110
Publication

21 Cerda Zolezzi, P.. "Ligaria cuneifolia flavonoid fractions modulate cell growth of normal lymphocytes and tumor cells as well as multidrug resistant cells", *Immunobiology*, 20050525
Publication

22 Giorgio Gaudiano, Tad H Koch, Mario Lo Bello, Marzia Nuccetelli et al. "Lack of glutathione conjugation to adriamycin in human breast cancer MCF-7/DOX cells", *Biochemical Pharmacology*, 2000
Publication

23 Putu Oka Samirana, Yosi Bayu Murti, Riris Istighfari Jenie, Erna Prawita Setyowati. "Antibacterial and cytotoxic activities of supernatant and mycelium extracts from fermentation of fungal symbiont *Trichoderma*

24

archiv.ub.uni-heidelberg.de

Internet Source

<1 %

25

www.jbc.org

Internet Source

<1 %

26

www.unej.ac.id

Internet Source

<1 %

27

123dok.com

Internet Source

<1 %

28

A. Lokhande, S. L. Ingale, S. H. Lee, S. Sen, C. Khong, B. J. Chae, I. K. Kwon. " Effects of dietary supplementation with (Merr.) on egg yolk cholesterol, excreta microflora and laying hen performance ", British Poultry Science, 2014

Publication

<1 %

29

Arisawa, Mitsuhiro, Yayoi Kasaya, Tohru Obata, Takuma Sasaki, Tomonori Nakamura, Takuya Araki, Koujiro Yamamoto, Akito Sasaki, Akihito Yamano, Mika Ito, Hiroshi Abe, Yoshihiro Ito, and Satoshi Shuto. "Design and Synthesis of Indomethacin Analogues That Inhibit P-Glycoprotein and/or Multidrug Resistant Protein without Cox Inhibitory

<1 %

Activity", Journal of Medicinal Chemistry,
2012.

Publication

30

E Fontaine. "Effects of permeability transition inhibition and decrease in cytochrome c content on doxorubicin toxicity in K562 cells", *Oncogene*, 04/27/2006

Publication

<1 %

31

Hamdoon A. Mohammed, Sulaiman A. Almahmoud, Minhajul Arfeen, Ashish Srivastava et al. "Phytochemical profiling, molecular docking, and anti-hepatocellular carcinoid bioactivity of extracts", *Arabian Journal of Chemistry*, 2022

Publication

<1 %

32

Sugihara, N.. "Effect of benzo[a]pyrene on P-glycoprotein-mediated transport in Caco-2 cell monolayer", *Toxicology*, 20060601

Publication

<1 %

33

Takeba, Yuko, Susumu Sekine, Toshio Kumai, Naoki Matsumoto, Sachiko Nakaya, Yashimitsu Tsuzuki, Yohei Yanagida, Hiroshi Nakano, Takeshi Asakura, Takehito Ohtsubo, and Shinichi Kobayashi. "Irinotecan-Induced Apoptosis Is Inhibited by Increased P-Glycoprotein Expression and Decreased p53 in Human Hepatocellular Carcinoma Cells", *Biological & Pharmaceutical Bulletin*, 2007.

Publication

<1 %

34 Vimal A. Patel, Michael J. Dunn, Andrey Sorokin. "Regulation of MDR-1 (P-glycoprotein) by Cyclooxygenase-2", Journal of Biological Chemistry, 2002

Publication

<1 %

35 Walker, J.. "Inhibition of P-glycoprotein function by XR9576 in a solid tumour model can restore anticancer drug efficacy", European Journal of Cancer, 200403

Publication

<1 %

36 japsonline.com

Internet Source

<1 %

37 journalarticle.ukm.my

Internet Source

<1 %

38 livrepository.liverpool.ac.uk

Internet Source

<1 %

39 www.jove.com

Internet Source

<1 %

40 Clifford W. Sachs, Ahmad R. Safa, Steadman D. Harrison, Robert L. Fine. "Partial Inhibition of Multidrug Resistance by Safingol Is Independent of Modulation of P-glycoprotein Substrate Activities and Correlated with Inhibition of Protein Kinase C", Journal of Biological Chemistry, 1995

Publication

<1 %

- | | | |
|----|---|------|
| 41 | Jin, J.. "Metabolites of ginsenosides as novel BCRP inhibitors", Biochemical and Biophysical Research Communications, 20060714
Publication | <1 % |
| 42 | Linda S. Steelman, William H. Chappell, Shaw M. Akula, Stephen L. Abrams et al.
"Therapeutic resistance in breast cancer cells can result from deregulated EGFR signaling", Advances in Biological Regulation, 2020
Publication | <1 % |
| 43 | Soo Yeon Chung. "Potent modulation of P-glycoprotein activity by naturally occurring phenylbutenoids from <i>Zingiber cassumunar</i> ", Phytotherapy Research, 04/2009
Publication | <1 % |
| 44 | Szakács, Gergely, Matthew D. Hall, Michael M. Gottesman et al. "Targeting the Achilles Heel of Multidrug-Resistant Cancer by Exploiting the Fitness Cost of Resistance", Chemical Reviews
Publication | <1 % |
| 45 | Submitted to University of Macau
Student Paper | <1 % |
| 46 | Vinay Kumar Gupta, Yashika Bhalla, Vikas Jaitak. "Impact of ABC transporters, glutathione conjugates in MDR and their | <1 % |

modulation by flavonoids: an overview", Medicinal Chemistry Research, 2013

Publication

47	aac.asm.org Internet Source	<1 %
48	coek.info Internet Source	<1 %
49	docslide.us Internet Source	<1 %
50	dspace.cuni.cz Internet Source	<1 %
51	es.scribd.com Internet Source	<1 %
52	he01.tci-thaijo.org Internet Source	<1 %
53	media.neliti.com Internet Source	<1 %
54	researchbank.rmit.edu.au Internet Source	<1 %
55	www.uni-heidelberg.de Internet Source	<1 %
56	Gudrun H. U. Paulsen, Trond Strickert, Anne B. L. Marthinsen, Steinar Lundgren. "Changes in Radiation Sensitivity and Steroid Receptor Content Induced by Hormonal Agents and	<1 %

Ionizing Radiation in Breast Cancer Cells in Vitro", Acta Oncologica, 2009

Publication

57

M.-H Teiten, L Bezdetnaya, J.-L Merlin, C Bour-Dill, M.E Pauly, M Dicato, F Guillemin. "Effect of meta-tetra(hydroxyphenyl)chlorin (mTHPC)-mediated photodynamic therapy on sensitive and multidrug-resistant human breast cancer cells", Journal of Photochemistry and Photobiology B: Biology, 2001

Publication

<1 %

58

Tan, Hui-Li, Kok-Gan Chan, Priyia Pusparajah, Learn-Han Lee, and Bey-Hing Goh. "Gynura procumbens: An Overview of the Biological Activities", Frontiers in Pharmacology, 2016.

Publication

<1 %

59

Chen, C.. "Quercetin: A potential drug to reverse multidrug resistance", Life Sciences, 20100911

Publication

<1 %

60

Fei Shen, Shaoyou Chu, Aimee K. Bence, Barbara Bailey et al. "Quantitation of Doxorubicin Uptake, Efflux, and Modulation of Multidrug Resistance (MDR) in MDR Human Cancer Cells", Journal of Pharmacology and Experimental Therapeutics, 2008

Publication

<1 %

61 Guohua An. "Herbal Supplement-Based Interactions", Enzyme- and Transporter-Based Drug-Drug Interactions, 2010 $<1\%$
Publication

62 Resistance to Targeted Anti-Cancer Therapeutics, 2015. $<1\%$
Publication

63 Shuji Kitagawa. "Inhibition of P-glycoprotein function by tea catechins in KB-C2 cells", Journal of Pharmacy and Pharmacology, 08/01/2004 $<1\%$
Publication

64 Xiuhua Meng, Jiawei Li, Mimi Li, Hongjiang Wang, Bingru Ren, Jian Chen, Weilin Li. "Traditional uses, phytochemistry, pharmacology and toxicology of the genus Gynura (Compositae): A comprehensive review", Journal of Ethnopharmacology, 2021 $<1\%$
Publication

65 Yaprak Dönmez, Laila Akhmetova, Özlem Darcansoy İşeri, Meltem Demirel Kars, Ufuk Gündüz. "Effect of MDR modulators verapamil and promethazine on gene expression levels of MDR1 and MRP1 in doxorubicin-resistant MCF-7 cells", Cancer Chemotherapy and Pharmacology, 2010 $<1\%$
Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On