

Research Article

IN VITRO RELEASE MODELING OF ASPIRIN FLOATING TABLETS USING DDSOLVER

Agus Siswanto^{1,2}, Achmad Fudholi^{1*}, Akhmad Kharis Nugroho¹, Sudibyo Martono¹

¹Faculty of Pharmacy,
Universitas Gadjah Mada
Sekip Utara Jogjakarta
55281, Indonesia

²Faculty of Pharmacy,
Muhammadiyah University
of Purwokerto, Jl. Raya
Dukuh Waluh Purwokerto
53142, Indonesia

Submitted: 14-01-2015

Revised: 10-02-2015

Accepted: 25-03-2015

*Corresponding author
Achmad Fudholi

Email:
fudholiugm@yahoo.com

ABSTRACT

Aspirin has low solubility in water therefore, dissolution is a rate limiting step for absorption. Floating tablet formulation is designed to improve the bioavailability of aspirin. The objective of this study was to determine *in vitro* dissolution study of aspirin floating tablet release kinetics model. The floating tablets were prepared by a direct compression method using Methocel K4M CR, NaHCO₃, Ethocel, Aerosil, and dicalcium phosphate anhydrous as excipients. Tablets were evaluated by different parameters such as physicochemical properties, floating lag time ($F_{lag\ time}$), total floating time, and dissolution. The result showed that the tablet mass has good flow properties of 13.54 g/sec. Aspirin floating tablets had a weight uniformity (CV=1.45%), good hardness (6.42kg), and low friability (0.158%). The tablet has a short $F_{lag\ time}$ of 25.16 sec and long floating time of 8 hours. Dissolution data were evaluated using DDSolver conducted by (1) Statistical parameters: $R^2_{adjusted}$, AIC, MSC; (2) Visual goodness of fit (GOF). The results showed that aspirin floating tablets release kinetics followed the Korsmeyer-Peppas model. Aspirin release occurs through the mechanism of anomalous transport which combines Fickian diffusion and polymer relaxation.

Key words: aspirin floating tablet, DDSolver, modeling of drug release



Temukan pesan, dokumen, foto, atau orang



Awal

Tulis

← Kembali ↩️ ⏪ ⏩ 📁 Arsipkan 📁 Pindahkan 🗑️ Hapus 🔒

📧 7 📄 ?



gus_ump@... 999+
agussiswanto1... 41

- ...
- Email Masuk 999+
- Belum Dibaca
- Berbintang
- Draft 154
- Terkirim
- Arsip
- Spam
- Sampah
- ^ Lebih sedikit

Tampilan Sembunyikan

- 📷 Foto
- 📄 Dokumen
- 📧 Langganan
- ✂️ Promo

Folder Sembunyikan

+ Folder Baru

• In Vitro Release Modeling of Aspirin Floating Tablets Using DDSolver AGUS Yahoo/Terkirim ★

AS • **agus siswanto** <gus_ump@...> Rab, 7 Jan 2015 jam 14.21 ★

Kepada:
mfi.ugm@gmail.com

Cc:
siswanto.kpp@gmail.com



In Vitro Rele... .doc
2.2MB

↩️ ⏪ ⏩ ...

Balas, Balas ke Semua atau Teruskan

Kirim **B** *I*

**Kode Voucher
Spesial Untukn**

**Asuransi Mobil
Dijamin Terhem**

All Risk Mulai
~~Rp 3.050.000/tahun~~
Rp 1.982.500/tahun

- 500+ Bengkel Rekanan
- Usta Mobil s.d 15 Tahun
- Klaim Mudah Semua Kerusakan

CEK SEKARANG

In Vitro Release Modeling of Aspirin Floating Tablets Using DDSolver

Agus Siswanto^{1,2}, Achmad Fudholi^{1*}, Akhmad Kharis Nugroho¹, Sudibyo Martono¹

¹Faculty of Pharmacy, Gadjah Mada University, Sekip Utara Jogjakarta
55281, Indonesia

²Faculty of Pharmacy, Muhammadiyah University of Purwokerto, Jl. Raya
Dukuh Waluh Purwokerto 53142, Indonesia

ABSTRACT

Aspirin has low solubility in water therefore, dissolution is a rate limiting step for absorption. Floating tablet formulation is designed to improve the bioavailability of aspirin. The objective of this study was to determine in vitro dissolution study of aspirin floating tablet release kinetics model. The floating tablets were prepared by a direct compression method using Methocel K4M CR, NaHCO₃, Ethocel, aerosil, and dicalcium phosphate anhydrous as excipients. Tablets were evaluated by different parameters such as fluidity, physicochemical properties, floating lag time ($F_{lag\ time}$), total floating time, and dissolution. The result showed that the tablet mass has good flow properties of 13.54 g/sec. Aspirin floating tablets had a weight uniformity (CV = 1.45%), good hardness (6.42 kg), and low friability (0.158%). The tablet has a short $F_{lag\ time}$ of 25.16 sec and long floating time of 8 hours. Dissolution data were evaluated using DDSolver conducted by (1) Statistical parameters: $R^2_{adjusted}$, AIC, MSC; (2) Visual goodness of fit (GOF). The results showed that aspirin floating tablets release kinetics followed the Korsmeyer-Peppas model. Aspirin release occurs through the mechanism of anomalous transport which combines Fickian diffusion and polymer relaxation.

Keywords: aspirin floating tablet, DDSolver, modeling of drug release

*Corresponding author, Hp. 08164220357
e-mail: fudholiugm@yahoo.com

INTRODUCTION

Besides analgesic, aspirin is also used as an antiplatelet for stroke prophylaxis (Laurer, 2002; Awtry & Loscalzo, 2000; Patrono & Rocca, 2008). Aspirin prevents blood from clotting by inhibiting cyclooxygenase, the enzyme responsible for the formation of thromboxane A₂, which is a mediator of platelet aggregating (Kannan *et al.*, 2010; Awtry & Loscalzo, 2000).

Aspirin has a low bioavailability due to first-pass effect and hydrolysis to salicylate metabolism in the gut wall (Sweetman, 2009). Aspirin is rapidly absorbed in the upper gastrointestinal tract, especially in the portion of the small intestine (Awtry & Loscalzo, 2000; Sweetman, 2009). Therefore, formulation of floating drug delivery system (floating on the gastric fluid) are designed to improve the bioavailability of aspirin (Suratri, 2008).

Aspirin has low solubility in water (1:300) (Moffat *et al.*, 2011; Sweetman, 2009) therefore, dissolution is one of the rate limiting step in their absorption and bioavailability (Gordon *et al.*, 1994; Wells, 2002). Information on the mechanisms and kinetics of dissolution is important to estimate the absorption of aspirin floating tablet. Several mathematical models were developed to analyze dissolution data such as the zero-order, first-order, Higuchi, Weibull, Korsmeyer-Peppas, Hixson-Crowell, Bakar-Lonsdale, and Hopfenberg. The nonlinear fitting of dissolution data can be performed using a professional statistical programs such as Micro-

Math Scientist, GraphPad Prism, SigmaPlot, or SYSTAT. However, these programs require the user to define the equations manually and need the initial value for each parameter (Zhang *et al.*, 2010). DDSolver is a new software developed for the kinetic analysis of dissolution data (Murtaza *et al.*, 2012) with a non-linear regression approach. DDSolver is a menu-driven add-in program for Microsoft Excel written in Visual Basic for Applications. Calculation using Excel offers a number of advantages over other software. The most attractive of advantages is the ease of use. Most scientists are already familiar with Excel because of its wide availability and high flexibility (Zhang *et al.*, 2010). This study was aimed to describe the dissolution data of aspirin floating tablet using DDSolver.

MATERIAL AND METHODS

Aspirin was obtained from Jilin Pharmaceutical Co. (Jilin, China), sodium bicarbonate from Tosoh Corporation (Tokyo, Japan), dicalcium phosphate anhydrous from Budenheim (Budenheim, Germany), Methocel K4M CR and Ethocel Premium Standard 20 from Colorcon Asia Pvt. Ltd. Hydrochloric acid and sodium chloride (analytical grade) were purchased from Merck (Darmstadt, Germany). All other chemicals used were of pharmaceutical grade.

Preparation of floating tablet

Tablets were prepared by direct compression method based on formulation in Table 1. Aspirin, Methocel K4M CR, sodium bicarbonate,

Ethocel, dicalcium phosphate anhydrous, and aerosil were mixed in a cube mixer at 30 rpm for 3 minutes. The tablet compression process used single punch tablet machine (Korsch, Germany). The weight of tablets were arranged at 200 mg and the hardness of tablet were controlled at 6-8 kg (Monsanto hardness tester).

Fluidity of tablet mass

The flow rate of tablet mass was determined by using flowability tester (Erweka, Germany). The funnel was filled with 20 g of tablet mass and the time needed to flow was measured.

Physical properties of tablet

Weight variation: Twenty tablets were randomly selected from the batch and weighed individually to check for weight variation. Hardness test: Monsanto hardness tester was used to determine the tablet hardness. Friability: About 20 tablets were selected from each batch and evaluated for friability using an abrasive tester (ERWEKA, Germany).

Floating lag time ($F_{lag\ time}$)

Tablet was immersed in beaker containing 100 mL of simulated gastric fluid (SGF) without pepsin pH 1.2. Time required for the tablet to rise to the surface and floating was determined as floating lag time (Patel *et al.*, 2007a).

In-vitro dissolution studies

The in vitro dissolution study was determined using Erweka DT 600 USP apparatus II (paddle method), using 900 mL SGF without pepsin pH 1.2 as dissolution medium at 37 ± 0.2 °C and 60 rpm (USP XXXII, 2008; Gopalakrishnan & Chentilnathan, 2011). Aliquots of 5 ml was taken out at intervals of 15, 30, 45, 60, 120, 180, 240, 300, 360, 420, and 480 minutes. Exactly 5 ml of fresh SGF was added to the dissolution vessel after each withdrawal to maintain a constant volume. The samples were analyzed by UV-Vis spectrophotometer (Genesys 10 Thermo) at 280 nm.

Determination of drug content

Twenty tablets were taken randomly from the batch, weighed, and powdered. A 200 mg of powder equivalent to 80 mg of aspirin was put into a 100 mL flask. Two mL of absolute ethanol and simulated gastric fluid (without pepsin) were added up to the mark. The solution was centrifuged for 60 minutes, then filtered with filter paper. The drug content was analyzed at the wavelength of aspirin (278 nm) and salicylic acid (304 nm) using a UV-Vis spectrophotometer. Drug content was then calculated with a simultaneous spectrophotometric method.

Data analysis

Physicochemical and floating properties were evaluated based on the applicable requirements. The dissolution data modeling was done using DDSolver. Evaluation of release kinetics model was conducted by (1)

Statistical parameters: R^2_{adjusted} , Akaike Information Criterion (AIC), Model Selection Criterion (MSC); and (2) Visual goodness of fit (GOF).

RESULTS AND DISCUSSION

The physicochemical properties of aspirin floating tablets

Aspirin floating tablets were prepared by a direct compression method. To obtain a good weight uniformity, the mass of tablets should have good flow properties. In this study, the mass of tablet has good flow properties with flow rate of 13.54 ± 0.46 g/sec (Fudholi, 1983). When the mass was compressed, it will flow properly at the compression chamber and produce the uniform weight and content.

Aspirin floating tablets have good physicochemical properties (Table 2). The tablets have a uniform weight with CV = 1.45 % so the active substance in each tablet was uniform (Davies, 2009). Tablets have good hardness of 6.42 kg (Gordon *et al.*, 1990) and low friability (<1%) therefore, it withstand to mechanical shocks and not easily destroyed. The content of aspirin in floating tablets within the required USP XXXII (2008) range of 95-105%.

In vitro buoyancy of aspirin floating tablets

One indicators of good floating tablet quality is the tablet's ability to float in the medium indicated by $F_{\text{lag time}}$ and floating time. The duration of floating indicates the time the tablets float in the dissolution medium.

Aspirin floating tablets was able to float during dissolution test (8 hours), intact and not eroded. This indicates that the composition of excipients used in the formulation was able to maintain the floating ability and the integrity of the tablets. Ethocel, aerosil, and dicalcium phosphate anhydrous are hydrophobic excipients that play a role in maintaining the integrity of the tablets in the dissolution medium.

$F_{lag\ time}$ indicates the speed of tablet to float in the dissolution medium. $F_{lag\ time}$ test (Table 2) showed that aspirin floating tablets within the requirements of less than 180 seconds (Patel *et al.*, 2007a). The sooner the tablet floats, the better the floating quality. Buoyancy was determined by the density of tablet. If the density is less than 1, the tablets will float properly (Patel *et al.*, 2007a; Dave *et al.*, 2004). Aspirin floating tablet buoyancy is determined by the composition of Methocel K4M CR, NaHCO_3 , and Ethocel as excipients. When NaHCO_3 reacts with SGF, the CO_2 gas was produced. Gas trapped in a hydrophilic gel Methocel K4M CR thereby, decreasing the density of the tablets (Patel *et al.*, 2007b). Ethocel, which is hydrophobic, will maintain the integrity of tablet to float in a long time. Therefore, the combination of these three excipients produced a floating tablet with short $F_{lag-time}$ and long floating duration.

In vitro dissolution profiles of aspirin floating tablets

Aspirin floating tablet dissolution test was conducted according to USP XXXII (2008). Aspirin tablets were designed to float in the stomach for a long time. Drug release will occur in the gastric; therefore, the SGF

(without pepsin) pH 1.2 was used as dissolution test medium. The dissolution profile was made by plotting the percentage of drug dissolved versus time (Figure 1). The amount of aspirin dissolved in the 60 minutes is 22.73% within USP XXXII (2008) range of 20-55%. The amount of aspirin released after 8 hours was 84.85%.

Dissolution Study of Aspirin Floating Tablet Release using DDSolver

In vitro drug release data were fitted to kinetics models such as the zero-order, first-order, Higuchi, Weibull, Korsmeyer-Peppas, Hixson-Crowell, Bakar-Lonsdale, and Hopfenberg using DDSolver. The input of dissolution data into DDSolver include: (1) percentage of drug dissolved, (2) dissolution time (minutes). Modeling dissolution by DDSolver were estimated by nonlinear regression in accordance with the equation in table 3.

Analysis model of visual goodness of fit (GOF) conducted in two methods: (1) the correlation of Q_0 (amount of aspirin dissolved in the experimental results) versus Q_c (the predicted amount of aspirin dissolved in the medium) (Figure 2); (2) the correlation of residuals versus time (Figure 3) (Nugroho *et al.*, 2014).

Correlation analysis of Q_0 versus Q_c in Figure 2 showed that the Korsmeyer-Peppas model was the most appropriate model to explain the phenomenon of aspirin dissolution in floating tablet. Kormeyer-Peppas models provide the less deviation between Q_0 and Q_c , so it appears in

Figure 2 that the distribution of dissolution data (Q_o) beyond the curve of predicted dissolution data (Q_c). Modeling kinetics of dissolution in zero-order, first-order, Higuchi, Weibull, Hixson-Crowell, Bakar-Lonsdale, and Hopfenberg produced the more deviation of Q_o and Q_c . This is reinforced by the residual correlation analysis ($Q_o - Q_c$) versus time indicates that Korsmeyer-Peppas models do not form a specific pattern in Figure 3.

DDSolver provides a number of statistical criteria to evaluate the dissolution model included the correlation coefficient ($R_{\text{obs-pre}}$), the coefficient of determination (R_{sqr} , R^2 , or COD), the adjusted coefficient of determination ($R_{\text{sqr_adj}}$ or R^2_{adjusted}), mean square error (MSE), standard deviation residuals (MSE_root or $S_{y.x}$), SS, WSS, Akaike Information Criterion (AIC), and Model Selection Criterion (MSC). Among these parameters, R^2_{adjusted} , AIC, MSC are the most popular and widely used in the identification of dissolution data modeling (Zhang *et al.*, 2010).

R^2_{adjusted} is the most appropriate parameter for comparing dissolution models with different parameters compared to R^2 . R^2_{adjusted} value will decrease with the increasing of dissolution model parameters. The best model is the model with the highest value of R^2_{adjusted} (Zhang *et al.*, 2010). Based on R^2_{adjusted} in Table 4, Korsmeyer-Peppas is the most appropriate model for this study.

AIC parameters have been used for more than 35 years in the dissolution data analysis because it is simple and easy to use (Zhang *et al.*, 2010). The model with the smallest AIC value is the most precise

model (Motulsky & Christopoulos, 2003). The MSC is a modified reciprocal form of the AIC and has been normalized so that it is independent of the scaling of the data points. If we compared several models, the model with the highest value of the MSC is the best model. In general, the model was appropriate when the MSC value > 2 or 3 (Zhang *et al.*, 2010). Dissolution model parameters in Table 4 showed that the Korsmeyer-Peppas models is the best model to explain the phenomenon of aspirin floating tablet dissolution.

Dissolution mechanism of aspirin floating tablet by Korsmeyer-Peppas

Korsmeyer-Peppas dissolution models explain the exponential relationship of dissolved drug fraction versus time.

$$F = k_{KP} t^n$$

F= fraction of drug dissolved at time t, a = drug dissolution constant, t = dissolution time (minutes), and n = dissolution exponent (Costa & Lobo, 2001).

Dissolution exponent (n) can be used to explain the mechanisms involved in the dissolution (Kalam *et al.*, 2007). The analysis result of aspirin floating tablet dissolution with Korsmeyer-Peppas gave n value of 0.712 (Table 5). For cylindrical dosage form (tablets) then $0.45 < n < 1$ indicates that dissolution occurs through the mechanism of anomalous transport (Costa & Lobo, 2001; Colombo *et al.*, 2000). The release of

aspirin from floating tablet occurs through the combination of Fickian diffusion and polymer relaxation. The dissolution of aspirin was influenced by the excipients in the tablet. Methocel K4M CR is a hydrophilic polymer HPMC. When it contacted with SGF, Methocel K4M CR will expand to form a thick gel. NaHCO₃ reacts with dissolution medium (SGF) produces CO₂ which form pores in the polymer gel Methocel K4M CR. Ethocel, which is hydrophobic, will maintain the integrity of the tablets to withdraw the massive erosion. Therefore, the release of aspirin occurs through Fickian diffusion pass through hydrophilic gel. The dissolution of aspirin was also determined by the relaxation ability of Methocel K4M CR and Ethocel in the medium.

CONCLUSION

Based on the results it can be concluded that aspirin floating tablets fulfilled the requirements of good quality tablet. The dissolution data evaluated with DDSolver showed that the kinetics release of floating tablets followed the Korsmeyer-Peppas model through anomalous transport mechanism which combines Fickian diffusion and polymer relaxation.

REFERENCES

- Awtry, E.H., Loscalzo, J., 2000, Aspirin, *Circulation*, **101**, 1206-1218
Costa, P., Lobo, J.M.S., 2001, Review Modeling and comparison of dissolution profiles, *European Journal of Pharmaceutical Sciences*, **13**,123-133
Colombo, P., Santi, P., Bettini, R., Brazel, C.S., Peppas, N.A., 2000, Drug Release from Swelling-Controlled System, in Wise, D.L.,

- Handbook of Pharmaceutical Controlled Technology, Marcel Dekker Inc., New York. pp 183-205
- Dave, B.S., Amin, A.F., Patel, M.M., 2004, Gastroretentive Drug Delivery System of Ranitidine Hydrochloride: Formulation and In Vitro Evaluation, *AAPS PharmSciTech*, **5(2)**, 1-6
- Davies, P., 2009, Oral Solid Dosage Form, In Gibson, M., Pharmaceutical Preformulation and Formulation, Informa healthcare, New York. pp 389
- Fudholi, A., 1983, Methodology Formulation in Direct Compression, *Medika*, **7(9)**, 586-593
- Gopalakrishnan, S., Chenthilnathan, A., 2011, Floating Drug Delivery Systems: A Review, *Journal of Pharmaceutical Science and Technology*, **3(2)**, 548-554.
- Gordon, R.E., Rosanke, T.W., Fonner, D.E., Anderson, N.R., Banker, G.S., 1990, Granulation Technology and Tablet Characterization, in Lieberman, H.A., Lachman, L., Schwartz, J.B., *Pharmaceutical Dosage Form: Tablet*, 2nd Ed., Vol 2, Marcel Dekker Inc., New York. pp 327-332
- Gordon, M. S., Ellis, D. J., Molony, B., 1994, In Vitro Dissolution Versus In Vivo Evaluation of Four Different Aspirin Products, *Drug Dev. Ind. Pharm.*, **20(10)**, 1711-1723
- Kannan, S., Manivannan, R., Balasubramaniam, A., and Kumar, N.S., 2010, Formulation and Evaluation of Aspirin Delayed Release Tablet, *IJCP*, 01, accessed 12.02.12, <<http://www.pharmacie-globale.info>>
- Kalam, M.A., Humayun, M., Parvez, N., Yadaz, S., Garg, A., Amin, S., Sultana, Y., Ali, A., 2007, Release Kinetics of Modified Pharmaceutical Dosage Forms: A Review, *J.Pharmaceutical Sciences*, **1**, 30-35
- Lauer, M.S., 2002, Aspirin For Primary Prevention of Coronary Events, *N Engl J Med*, **346(19)**, 1468-1474
- Moffat, A.C., Osselton, M.D., Widdop, B., and Watts, J., 2011, Clarke's Analysis of Drugs and Poisons in Pharmaceuticals, Body Fluids and Postmortem Material, fourth edition, Pharmaceutical Press, London-Chicago. pp 924-926
- Motulsky, H.J., Christopoulos, A., 2003, Fitting Models To Biological Data Using Linear and Nonlinear Regression. A Practical Guide To Curve Fitting, GraphPad Software, Inc., San Diego CA. pp 143-148
- Murtaza, G., Ahmad, M., Khan, S.A., Hussain, I., 2012, Evaluation of Cefixime-Loaded Chitosan Microspheres: Analysis of Dissolution Data Using DDSolver, *Dissolution Technologies*, 13-19
- Nugroho, A.K., Binnarjo, A., Hakim, A.R., Ermawati, Y., 2014, Compartmental Modeling Approach of Losartan Transdermal Transport In Vitro, *Indonesian J. Pharm.*, Vol. 25 No. 1, 31-38

- Patel D.M., Patel N.M., Pandya N.N., Jogani, P.D., 2007a, Gastroretentive Drug Delivery System of Carbamazepine: Formulation Optimization Using Simplex Lattice Design: A Technical Note, *AAPS PharmSciTech*, **8(1)**, E1:E5, accessed 24.10.11, <<http://www.aapspharmstech.org>>
- Patel, V.F., Patel, N.M., 2007b, Statistical Evaluation of Influence of Viscosity and Content of Polymer on Dipyridamole Release From Floating Matrix Tablets: A Technical Note, *AAPS PharmSciTech*, **8(3)**, E2-E5, accessed 24.10.11, <<http://www.aapspharmstech.org>>
- Patrono, C., Rocca, B., 2008, Aspirin: Promise and Resistance in the New Millennium, *Arterioscler Thromb Vasc Biol*, **28**, s25-s32, accessed 16.11.11, <<http://atvb.ahajournals.org>>.
- Suratri, A., 2008, Floating Drug Delivery Systems To Increase Gastric Retention of Drugs: A Review, *RJPT*, **1(4)**, accessed 02.04.11, <<http://www.rjptonline.org/>>.
- Sweetman, S. C., 2009, Martindale The Complete Drug Reference, Thirty-sixth edition, Pharmaceutical Press, London. pp 20-25
- USP, 2008, United States Pharmacopeial XXXII, accessed 17.02.12, <<http://www.pharmacopeia.cn/>>.
- Wells, J., 2002, Pharmaceutical preformulation: The Physicochemical Properties of Drug Substances, in Aulton, M.E., *Pharmaceutics: The Science of Dosage Form Design*, Ed. II, Churchill Livingstone, Edinburg. pp 113-138
- Zhang, Y., Huo, M., Zhou, J., Zou, A., Li, W., Yao, C., Xie, S., 2010, DDSolver: An Add-In Program for Modeling and Comparison of Drug Dissolution Profiles, *The AAPS Journal*, **Vol. 12 No. 3**, 263-271

Table 1. Formulation of aspirin floating tablet

Ingredient	Amount (mg) / tablet
Aspirin	80.00
Methocel K4M CR	33.87
NaHCO ₃	30.37
Ethocel	35.76
Aerosil	2.00
Dicalcium phosphate anhydrous	18.00
Total weight (mg)	200.0

Table 2. Physicochemical properties of aspirin floating tablets

Parameters	Mean ± SD
Weight of tablet (mg)	201.1 ± 0.6
Friability (%)	0.16
Hardness (kg)	6.42 ± 0.58
F _{lag} time (sec)	25.16 ± 4.60
Total floating time (hours)	> 8
Drug content (aspirin) (%)	99.55 ± 1.99
Drug content (salysilic acid) (%)	1.24 ± 0.09

Table 3. Models Available in DDSolver for Fitting Drug Release Data

Model	Equation
Zero order	$F = k_0 t$
First order	$F = 100 [1 - \text{Exp}(-k_1 t)]$
Higuchi	$F = k_H t^{0.5}$
Hixson-Crowell	$F = 100 [1 - (1 - k_{HC} t)^3]$
Hopfenberg	$F = 100 [1 - (1 - k_{HB} t)^n]$
Weibull	$F = 100 \{1 - \text{Exp}[-(t^\beta)/\alpha]\}$
Korsmeyer-Peppas	$F = k_{KP} t^n$
Baker-Lonsdale	$3/2 [1 - (1 - F/100)^{(2/3)}] - F/100 = k_{BL} t$

Note: F = drug released (%); t = time (minutes)

Table 4. Statistical parameters of models to describe release of aspirin floating tablets

Model	Statistical parameters of models (mean±SD)		
	R ² _{adjusted}	AIC	MSC
Zero order	0.938±0.037	75.38±8.65	2.62±0.77
First order	0.982±0.010	61.35±5.98	3.79±0.51
Higuchi	0.951±0.027	73.42±7.22	2.79±0.55
Korsmeyer-Peppas	0.994±0.002	48.86±4.80	4.83±0.46
Hixson-Crowell	0.983±0.009	60.26±7.91	3.88±0.75
Hopfenberg	0.988±0.007	57.81±6.22	4.09±0.60
Baker-Lonsdale	0.910±0.033	81.67±4.91	2.10±0.35
Weibull	0.985±0.009	59.94±6.38	3.91±0.55

Table 5. Dissolution parameters of aspirin floating tablets by Korsmeyer-Peppas

Parameter	Mean ± SD
k _{KP}	1.175 ± 0.562
n	0.712 ± 0.091




Figure 1. Drug release of aspirin floating tablets




Figure 2. Observed data of drug release of aspirin floating tablets in dissolution studies, presented together with the model prediction using zero order, first order, Higuchi, Hixson-Crowell, Hopfenberg, Baker-Lonsdale, Korsmeyer-Peppas, and Weibull




Figure 3. Goodness of fit (GOF) evaluations based on correlation of residual versus time indicating that Korsmeyer-Peppas models do not form a specific pattern



Temukan pesan, dokumen, foto, atau orang



Awal

Tulis

← Kembali ↩️ ↶️ ➡️ 📁 Arsipkan 📁 Pindahkan 🗑️ Hapus 🛡️

📧 7 📄 ? ⚙️

gus_ump@... 999+
agussiswanto1... 41

• Acknowledgement Manuscript Submission MS MFI-2015-02 Yahoo/Email M... ★

...
Email Masuk 999+
Belum Dibaca
Berbintang
Draft 154
Terakhir
Arsip
Spam
Sampah
^ Lebih sedikit

• mfi@ugm.ac.id Sen, 12 Jan 2015 jam 09.36 ★
Kepada: gus_ump@yahoo.com

Dear Mr. Agus Siswanto

This message is to acknowledge the receipt of the manuscript you submitted via email to the Indonesian Journal of Pharmacy. Your manuscript ID:MFI-2015-02.

Entitle : "In Vitro Release Modeling of Aspirin Floating Tablets Using DDSolver"

Your manuscript has been received. Your paper will be reviewed by three or more reviewers assigned by the editorial board and final decision made by the editor will be informed by this email.

You can monitor the review process for your paper by emailing us on the "Status of my manuscript No. MFI-2015-02".

Thank you very much for submitting your manuscript to the Indonesian Journal of Pharmacy.

Sincerely yours,

Prof. Dr. Abdul Rohman, M.Si., Apt
Associate editor
Indonesian Journal of Pharmacy

Info UGM: Sampaikan saran dan keluhan terhadap layanan UGM melalui sistem informasi aspirasi <http://aspirasi.ugm.ac.id>

↩️ ↶️ ➡️ ...

Tampilan Sembunyikan
Foto
Dokumen
Langganan
Promo
Folder Sembunyikan
+ Folder Baru

Balas, Balas ke Semua atau Teruskan



mfi@ugm.ac.id 🔍

mfi@ugm.ac.id
+ Tambah ke kontak

Belanja Sports Diskor
50K

istyle.id

Sports Favorit Diskon
50K

istyle.id



Temukan pesan, dokumen, foto, atau orang



Awal

Tulis

← Kembali ↶ ↷ ➡ Arsipkan 📁 Pindahkan 📁 Hapus ✖

📧 7 📄 ?



- gus_ump@...** 999+
- agussiswanto1... 41
- ...
- Email Masuk 999+
- Belum Dibaca
- Berbintang
- Draft 154
- Terkirim
- Arsip
- Spam
- Sampah
- ^ Lebih sedikit
- Tampilan Sembunyikan
- 📷 Foto
- 📄 Dokumen
- 📧 Langganan
- ✂️ Promo
- Folder Sembunyikan
- + Folder Baru




Review manuscript IJP-2015-2 Yahoo/Email M... ★



mfi ugm <mfi.ugm@gmail.com> Sel, 17 Mar 2015 jam 10.10
Kepada: gus_ump@yahoo.com

Dear Agus Siswanto
Please revise the review comment in attachment.

📄 Unduh semua lampiran sebagai file zip

- | | | |
|---|---|---|
|  |  |  |
| IJP-02_rev27... .pdf
535.1kB | IJP-02-Evalua... .pdf
385.2kB | IJP-02_rev2... .docx
1.2MB |



Balas, Balas ke Semua atau Teruskan

Kirim 📎 📄 📧 😊 🔗 B I 🗑️



mfi.ugm@gmail.com 🔍
mfi.ugm@gmail.com
+ Tambah ke kontak

Asuransi Mobil Dijamin Terher

All Risk Mulai
~~Rp 3.050.000/tahun~~
Rp 1.982.500/tahun

- 500+ Bengkel Rekanan
- Usia Mobil s.d 15 Tahun
- Klaim Mudah Semua Kerusakan

CEK SEKARANG

Asuransi Mobil Dijamin Terher

All Risk Mulai
~~Rp 3.050.000/tahun~~
Rp 1.982.500/tahun

- 500+ Bengkel Rekanan
- Usia Mobil s.d 15 Tahun
- Klaim Mudah Semua Kerusakan

CEK SEKARANG

In Vitro Release Modeling of Aspirin Floating Tablets Using DDSolver

ABSTRACT

Aspirin has low solubility in water therefore, dissolution is a rate limiting step for absorption. Floating tablet formulation is designed to improve the bioavailability of aspirin. The objective of this study was to determine in vitro dissolution study of aspirin floating tablet release kinetics model. The floating tablets were prepared by a direct compression method using Methocel K4M CR, NaHCO₃, Ethocel, aerosil, and dicalcium phosphate anhydrous as excipients. Tablets were evaluated by different parameters such as fluidity (not parameter of tablet, its parameter of tablet mass?), physicochemical properties, floating lag time ($F_{lag\ time}$), total floating time, and dissolution. The result showed that the tablet mass has good flow properties of 13.54 g/sec. Aspirin floating tablets had a weight uniformity (CV = 1.45%), good hardness (6.42 kg), and low friability (0.158%). The tablet has a short $F_{lag\ time}$ of 25.16 sec and long floating time of 8 hours. Dissolution data were evaluated using DDSolver conducted by (1) Statistical parameters: $R^2_{adjusted}$, AIC, MSC; (2) Visual goodness of fit (GOF). The results showed that aspirin floating tablets release kinetics followed the Korsmeyer-Peppas model. Aspirin release occurs through the mechanism of anomalous transport which combines Fickian diffusion and polymer relaxation.

Keywords: aspirin floating tablet, DDSolver, modeling of drug release

Formatted: Font color: Red

INTRODUCTION

Besides analgesic, aspirin is also used as an analgesic and antiplatelet for stroke prophylaxis (Laurer, 2002; Awtry & Loscalzo, 2000; Patrono & Rocca, 2008). Aspirin prevents blood from clotting by inhibiting cyclooxygenase, the enzyme responsible for the formation of thromboxane A₂, which is a mediator of platelet aggregating (Kannan *et al.*, 2010; Awtry & Loscalzo, 2000).

Aspirin has a low bioavailability due to first-pass effect and hydrolysis to salicylate metabolism in the gut wall (Sweetman, 2009). Aspirin is rapidly absorbed in the upper gastrointestinal tract, especially in the portion of the small intestine (Awtry & Loscalzo, 2000; Sweetman, 2009). Therefore, formulation of floating drug delivery system (floating on the gastric fluid) are designed to improve the bioavailability of aspirin (Suratri, 2008).

Aspirin has low solubility in water (1:300) (Moffat *et al.*, 2011; Sweetman, 2009) therefore, dissolution is one of the rate limiting step in their absorption and bioavailability (Gordon *et al.*, 1994; Wells, 2002). Information on the mechanisms and kinetics of dissolution is important to estimate the absorption of aspirin floating tablet. Several mathematical models were developed to analyze dissolution data such as the zero-order, first-order, Higuchi, Weibull, Korsmeyer-Peppas, Hixson-Crowell, Bakar-Lonsdale, and Hopfenberg. The nonlinear fitting of dissolution data can be performed using a professional statistical programs such as Micro-Math Scientist, GraphPad Prism, SigmaPlot, or SYSTAT. However, these

programs require the user to define the equations manually and need the initial value for each parameter (Zhang *et al.*, 2010). DDSolver is a new software developed for the kinetic analysis of dissolution data (Murtaza *et al.*, 2012) with a non-linear regression approach. DDSolver is a menu-driven add-in program for Microsoft Excel written in Visual Basic for Applications. Calculation using Excel offers a number of advantages over other software. The most attractive of advantages is the ease of use. Most scientists are already familiar with Excel because of its wide availability and high flexibility (Zhang *et al.*, 2010). This study was aimed to describe the dissolution data of aspirin floating tablet using DDSolver.

MATERIAL AND METHODS

Aspirin was obtained from Jilin Pharmaceutical Co. (Jilin, China), sodium bicarbonate from Tosoh Corporation (Tokyo, Japan), dicalcium phosphate anhydrous from Budenheim (Budenheim, Germany), Methocel K4M CR and Ethocel Premium Standard 20 from Colorcon Asia Pvt. Ltd. Hydrochloric acid and sodium chloride (analytical grade) were purchased from Merck (Darmstadt, Germany). All other chemicals used were of pharmaceutical grade.

Preparation of floating tablet

Tablets were prepared by direct compression method based on formulation in Table 1. Aspirin, Methocel K4M CR, sodium bicarbonate, Ethocel, dicalcium phosphate anhydrous, and aerosil were mixed in a cube

mixer at 30 rpm for 3 minutes. The tablet compression process used single punch tablet machine (Korsch, Germany). The weight of tablets were arranged at 200 mg and the hardness of tablet were controlled at 6-8 kg (Monsanto hardness tester).

Fluidity of tablet mass

The flow rate of tablet mass was determined by using flowability tester (Erweka, Germany). The funnel was filled with 20 g of tablet mass and the time needed to flow was measured.

Physical properties of tablet

Weight variation: Twenty tablets were randomly selected from the batch and weighed individually to check for weight variation. Hardness test: Monsanto hardness tester was used to determine the tablet hardness. Friability: About 20 tablets were selected from each batch and evaluated for friability using an abrasive tester (ERWEKA, Germany).

Floating lag time ($F_{lag\ time}$)

Tablet was immersed in beaker containing 100 mL of simulated gastric fluid (SGF) without pepsin pH 1.2. Time required for the tablet to rise to the surface and floating was determined as floating lag time (Patel *et al.*, 2007a).

In-vitro dissolution studies

The in vitro dissolution study was determined using Erweka DT 600 USP apparatus II (paddle method), using 900 mL SGF without pepsin pH 1.2 as dissolution medium at 37 ± 0.2 °C and 60 rpm (USP XXXII, 2008; Gopalakrishnan & Chentilnathan, 2011). Aliquots of 5 ml was taken out at intervals of 15, 30, 45, 60, 120, 180, 240, 300, 360, 420, and 480 minutes. Exactly 5 ml of fresh SGF was added to the dissolution vessel after each withdrawal to maintain a constant volume. The samples were analyzed by UV-Vis spectrophotometer (Genesys 10 Thermo) at 280 nm.

Determination of drug content

Twenty tablets were taken randomly from the batch, weighed, and powdered. A 200 mg of powder equivalent to 80 mg of aspirin was put into a 100 mL flask. Two mL of absolute ethanol and simulated gastric fluid (without pepsin) were added up to the mark. The solution was centrifuged for 60 minutes, then filtered with filter paper. The drug content was analyzed at the wavelength of aspirin (278 nm) and salicylic acid (304 nm) using a UV-Vis spectrophotometer. Drug content was then calculated with a simultaneous spectrophotometric method.

Data analysis

Physicochemical and floating properties were evaluated based on the applicable requirements. The dissolution data modeling was ~~done~~ calculated using DDSolver. Evaluation of release kinetics model was

Formatted: Font color: Text 2, Strikethrough

conducted by (1) Statistical parameters: R^2_{adjusted} , Akaike Information Criterion (AIC), Model Selection Criterion (MSC); and (2) Visual goodness of fit (GOF).

RESULTS AND DISCUSSION

The physicochemical properties of aspirin floating tablets

Aspirin floating tablets were prepared by a direct compression method. To obtain a good weight uniformity, the mass of tablets should have good flow properties. In this study, the mass of tablet has good flow properties with flow rate of 13.54 ± 0.46 g/sec (Fudholi, 1983). When the mass was compressed, it will flow properly at the compression chamber and produce the uniform weight and content.

Aspirin floating tablets have good physicochemical properties (Table 2). The tablets have a uniform weight with CV = 1.45 % so the active substance in each tablet was uniform (Davies, 2009). Tablets have good hardness of 6.42 kg (Gordon *et al.*, 1990) and low friability (<1%) therefore, it withstand to mechanical shocks and not easily destroyed. The content of aspirin in floating tablets within the required USP XXXII (2008) range of 95-105%.

In vitro buoyancy of aspirin floating tablets

One ~~indicators parameter~~ of good floating tablet quality is the tablet's ability to float in the medium indicated by $F_{\text{lag time}}$ and floating time. The

Formatted: Font color: Text 2, Strikethrough

Formatted: Font color: Red

duration of floating indicates the time ~~the of~~ tablets float in the dissolution medium. Aspirin floating tablets was able to float during dissolution test (8 hours), intact and not eroded. ~~(please write the requirement of the floating time of the tablet so we can say the tablet has good parameter of this character)~~ This indicates that the composition of excipients used in the formulation was able to maintain the floating ability and the integrity of the tablets. Ethocel, aerosil, and dicalcium phosphate anhydrous are hydrophobic excipients that play a role in ~~the~~ maintaining ~~the of~~ integrity of the tablets in the dissolution medium.

Formatted: Font color: Text 2, Strikethrough

Formatted: Font color: Text 2, Strikethrough

Formatted: Font color: Red

$F_{lag\ time}$ indicates the speed of tablet to float in the dissolution medium. $F_{lag\ time}$ test (Table 2) showed that aspirin floating tablets within the requirements of less than 180 seconds (Patel *et al.*, 2007a). The sooner the tablet floats, the better the floating quality. Buoyancy was determined by the density of tablet. If the density is less than 1, the tablets will float properly (Patel *et al.*, 2007a; Dave *et al.*, 2004). Aspirin floating tablet buoyancy is determined by the composition of Methocel K4M CR, NaHCO_3 , and Ethocel as excipients. When NaHCO_3 reacts with SGF, the CO_2 gas was produced. Gas trapped in a hydrophilic gel Methocel K4M CR thereby, decreasing the density of the tablets (Patel *et al.*, 2007b). Ethocel, which is hydrophobic, will maintain the integrity of tablet to float in a long time. Therefore, the combination of these three excipients produced a floating tablet with short $F_{lag-time}$ and long floating duration.

In vitro dissolution profiles of aspirin floating tablets

Aspirin floating tablet dissolution test was conducted according to USP XXXII (2008). Aspirin tablets were designed to float in the stomach for a long time. Drug release will occur in the gastric; therefore, the SGF (without pepsin) pH 1.2 was used as dissolution test medium. The dissolution profile was made by plotting the percentage of drug dissolved versus time (Figure 1). The amount of aspirin dissolved in the 60 minutes is 22.73% within USP XXXII (2008) range of 20-55%. The amount of aspirin released after 8 hours was 84.85%.

Dissolution Study of Aspirin Floating Tablet Release using DDSolver

In vitro drug release data were fitted to kinetics models such as the zero-order, first-order, Higuchi, Weibull, Korsmeyer-Peppas, Hixson-Crowell, Bakar-Lonsdale, and Hopfenberg using DDSolver. The input of dissolution data into DDSolver include: (1) percentage of drug dissolved, (2) dissolution time (minutes). Modeling dissolution by DDSolver were estimated by nonlinear regression in accordance with the equation in table 3.

Analysis model of visual goodness of fit (GOF) conducted in two methods: (1) the correlation of Q_o (amount of aspirin dissolved in the experimental results) versus Q_c (the predicted amount of aspirin dissolved in the medium) (Figure 2); (2) the correlation of residuals versus time (Figure 3) (Nugroho *et al.*, 2014).

Correlation analysis of Q_o versus Q_c in Figure 2 showed that the Korsmeyer-Peppas model was the most appropriate model to explain the phenomenon of aspirin dissolution in floating tablet. Korsmeyer-Peppas models provide the less deviation between Q_o and Q_c , so it appears in Figure 2 that the distribution of dissolution data (Q_o) beyond the curve of predicted dissolution data (Q_c). Modeling kinetics of dissolution in zero-order, first-order, Higuchi, Weibull, Hixson-Crowell, Bakar-Lonsdale, and Hopfenberg produced the more deviation of Q_o and Q_c . This is reinforced by the residual correlation analysis ($Q_o - Q_c$) versus time indicates that Korsmeyer-Peppas models do not form a specific pattern in Figure 3.

DDSolver provides a number of statistical criteria to evaluate the dissolution model included the correlation coefficient ($R_{obs-pre}$), the coefficient of determination ($Rsqr$, R^2 , or COD), the adjusted coefficient of determination ($Rsqr_{adj}$ or $R^2_{adjusted}$), mean square error (MSE), standard deviation residuals (MSE_{root} or $Sy.x$), SS, WSS, Akaike Information Criterion (AIC), and Model Selection Criterion (MSC). Among these parameters, $R^2_{adjusted}$, AIC, MSC are the most popular and widely used in the identification of dissolution data modeling (Zhang *et al.*, 2010).

$R^2_{adjusted}$ is the most appropriate parameter for comparing dissolution models with different parameters compared to R^2 . $R^2_{adjusted}$ value will decrease with the increasing of dissolution model parameters. The best model is the model with the highest value of $R^2_{adjusted}$ (Zhang *et al.*, 2010).

Based on R^2_{adjusted} in Table 4, Korsmeyer-Peppas is the most appropriate model for this study.

AIC parameters have been used for more than 35 years in the dissolution data analysis because it is simple and easy to use (Zhang *et al.*, 2010). The model with the smallest AIC value is the most precise model (Motulsky & Christopoulos, 2003). The MSC is a modified reciprocal form of the AIC and has been normalized so that it is independent of the scaling of the data points. If we compared several models, the model with the highest value of the MSC is the best model. In general, the model was appropriate when the MSC value > 2 or 3 (Zhang *et al.*, 2010). Dissolution model parameters in Table 4 showed that the Korsmeyer-Peppas models is the best model to explain the phenomenon of aspirin floating tablet dissolution.

Dissolution mechanism of aspirin floating tablet by Korsmeyer-Peppas

Korsmeyer-Peppas dissolution models explain the exponential relationship of dissolved drug fraction versus time.

$$F = k_{KP} t^n$$

F= fraction of drug dissolved at time t, a = drug dissolution constant, t = dissolution time (minutes), and n = dissolution exponent (Costa & Lobo, 2001).

Dissolution exponent (n) can be used to explain the mechanisms involved in the dissolution (Kalam *et al.*, 2007). The analysis result of aspirin

floating tablet dissolution with Korsmeyer-Peppas gave n value of 0.712 (Table 5). For cylindrical dosage form (tablets) then $0.45 < n < 1$ indicates that dissolution occurs through the mechanism of anomalous transport (Costa & Lobo, 2001; Colombo *et al.*, 2000). The release of aspirin from floating tablet occurs through the combination of Fickian diffusion and polymer relaxation. The dissolution of aspirin was influenced by the excipients in the tablet. Methocel K4M CR is a hydrophilic polymer HPMC. When it contacted with SGF, Methocel K4M CR will expand to form a thick gel. NaHCO_3 reacts with dissolution medium (SGF) produces CO_2 which form pores in the polymer gel Methocel K4M CR. Ethocel, which is hydrophobic, will maintain the integrity of the tablets to withdraw the massive erosion. Therefore, the release of aspirin occurs through Fickian diffusion pass through hydrophilic gel. The dissolution of aspirin was also determined by the relaxation ability of Methocel K4M CR and Ethocel in the medium.

CONCLUSION

Based on the results it can be concluded that aspirin floating tablets fulfilled the requirements of good quality tablet. The dissolution data evaluated with DDSolver showed that the kinetics release of floating tablets followed the Korsmeyer-Peppas model through anomalous transport mechanism which combines Fickian diffusion and polymer relaxation.

REFERENCES

Awtry, E.H., Loscalzo, J., 2000, Aspirin, *Circulation*, **101**, 1206-1218

- Costa, P., Lobo, J.M.S., 2001, Review Modeling and comparison of dissolution profiles, *European Journal of Pharmaceutical Sciences*, **13**,123-133
- Colombo, P., Santi, P., Bettini, R., Brazel, C.S., Peppas, N.A., 2000, Drug Release from Swelling-Controlled System, in Wise, D.L., Handbook of Pharmaceutical Controlled Technology, Marcel Dekker Inc., New York. pp 183-205
- Dave, B.S., Amin, A.F., Patel, M.M., 2004, Gastroretentive Drug Delivery System of Ranitidine Hydrochloride:Formulation and In Vitro Evaluation, *AAPS PharmSciTech*, **5(2)**, 1-6
- Davies, P., 2009, Oral Solid Dosage Form, In Gibson, M., Pharmaceutical Preformulation and Formulation, Informa healthcare, New York. pp 389
- Fudholi, A., 1983, Methodology Formulation in Direct Compression, *Medika*, **7(9)**, 586-593
- Gopalakrishnan, S., Chentilnathan, A., 2011, Floating Drug Delivery Systems: A Review, *Journal of Pharmaceutical Science and Technology*, **3(2)**, 548-554.
- Gordon, R.E., Rosanke, T.W., Fonner, D.E., Anderson, N.R., Banker, G.S., 1990, Granulation Technology and Tablet Characterization, in Lieberman, H.A., Lachman, L., Schwartz, J.B., Pharmaceutical Dosage Form: Tablet, 2nd Ed., Vol 2, Marcel Dekker Inc., New York. pp 327-332
- Gordon, M. S., Ellis, D. J., Molony, B., 1994, In Vitro Dissolution Versus In Vivo Evaluation of Four Different Aspirin Products, *Drug Dev. Ind. Pharm.*, **20(10)**, 1711-1723
- Kannan, S., Manivannan, R., Balasubramaniam, A., and Kumar, N.S., 2010, Formulation and Evaluation of Aspirin Delayed Release Tablet, *IJCP*, 01, accessed 12.02.12, <<http://www.pharmacie-globale.info>>
- Kalam, M.A., Humayun, M., Parvez, N., Yadaz, S., Garg, A., Amin, S., Sultana, Y., Ali, A., 2007, Release Kinetics of Modified Pharmaceutical Dosage Forms: A Review, *J.Pharmaceutical Sciences*, **1**, 30-35
- Lauer, M.S., 2002, Aspirin For Primary Prevention of Coronary Events, *N Engl J Med*, **346(19)**, 1468-1474
- Moffat, A.C., Osselton, M.D., Widdop, B., and Watts, J., 2011, Clarke's Analysis of Drugs and Poisons in Pharmaceuticals, Body Fluids and Postmortem Material, fourth edition, Pharmaceutical Press, London-Chicago. pp 924-926
- Motulsky, H.J., Christopoulos, A., 2003, Fitting Models To Biological Data Using Linear and Nonlinear Regression. A Practical Guide To Curve Fitting, GraphPad Software, Inc., San Diego CA. pp 143-148
- Murtaza, G., Ahmad, M., Khan, S.A., Hussain, I., 2012, Evaluation of Cefixime-Loaded Chitosan Microspheres: Analysis of Dissolution Data Using DDSolver, *Dissolution Technologies*, 13-19

- Nugroho, A.K., Binnarjo, A., Hakim, A.R., Ermawati, Y., 2014, Compartmental Modeling Approach of Losartan Transdermal Transport In Vitro, *Indonesian J. Pharm.*, Vol. 25 No. 1, 31-38
- Patel D.M., Patel N.M., Pandya N.N., Jogani, P.D., 2007a, Gastroretentive Drug Delivery System of Carbamazepine: Formulation Optimization Using Simplex Lattice Design: A Technical Note, *AAPS PharmSciTech*, **8(1)**, E1:E5, accessed 24.10.11, <<http://www.aapspharmscitech.org>>
- Patel, V.F., Patel, N.M., 2007b, Statistical Evaluation of Influence of Viscosity and Content of Polymer on Dipyridamole Release From Floating Matrix Tablets: A Technical Note, *AAPS PharmSciTech*, **8(3)**, E2-E5, accessed 24.10.11, <<http://www.aapspharmscitech.org>>
- Patrono, C., Rocca, B., 2008, Aspirin: Promise and Resistance in the New Millennium, *Arterioscler Thromb Vasc Biol*, **28**, s25-s32, accessed 16.11.11, <<http://atvb.ahajournals.org>>.
- Suratri, A., 2008, Floating Drug Delivery Systems To Increase Gastric Retention of Drugs: A Review, *RJPT*, **1(4)**, accessed 02.04.11, <<http://www.rjptonline.org/>>.
- Sweetman, S. C., 2009, Martindale The Complete Drug Reference, Thirty-sixth edition, Pharmaceutical Press, London. pp 20-25
- USP, 2008, United States Pharmacopeial XXXII, accessed 17.02.12, <<http://www.pharmacopeia.cn/>>.
- Wells, J., 2002, Pharmaceutical preformulation: The Physicochemical Properties of Drug Substances, in Aulton, M.E., *Pharmaceutics: The Science of Dosage Form Design*, Ed. II, Churchill Livingstone, Edinburg. pp 113-138
- Zhang, Y., Huo, M., Zhou, J., Zou, A., Li, W., Yao, C., Xie, S., 2010, DDSolver: An Add-In Program for Modeling and Comparison of Drug Dissolution Profiles, *The AAPS Journal*, **Vol. 12 No. 3**, 263-271

Table 1. Formulation of aspirin floating tablet

Ingredient	Amount (mg) / tablet
Aspirin	80.00
Methocel K4M CR	33.87
NaHCO ₃	30.37
Ethocel	35.76
Aerosil	2.00
Dicalcium phosphate anhydrous	18.00
Total weight (mg)	200.0

Table 2. Physicochemical properties of aspirin floating tablets

Parameters	Mean ± SD
Weight of tablet (mg)	201.1 ± 0.6
Friability (%)	0.16
Hardness (kg)	6.42 ± 0.58
F _{lag} time (sec)	25.16 ± 4.60
Total floating time (hours)	> 8
Drug content (aspirin) (%)	99.55 ± 1.99
Drug content (salicylic acid) (%)	1.24 ± 0.09

Table 3. Models Available in DDSolver for Fitting Drug Release Data

Model	Equation
Zero order	$F = k_0 t$
First order	$F = 100 [1 - \text{Exp}(-k_1 t)]$
Higuchi	$F = k_H t^{0.5}$
Hixson-Crowell	$F = 100 [1 - (1 - k_{HC} t)^3]$
Hopfenberg	$F = 100 [1 - (1 - k_{HB} t)^n]$
Weibull	$F = 100 \{1 - \text{Exp}[-(t^\beta)/\alpha]\}$
Korsmeyer-Peppas	$F = k_{KP} t^n$
Baker-Lonsdale	$3/2 [1 - (1 - F/100)^{2/3}] - F/100 = k_{BL} t$

Note: F = drug released (%); t = time (minutes)

Table 4. Statistical parameters of models to describe release of aspirin floating tablets

Model	Statistical parameters of models (mean±SD)		
	R ² _{adjusted}	AIC	MSC
Zero order	0.938±0.037	75.38±8.65	2.62±0.77
First order	0.982±0.010	61.35±5.98	3.79±0.51
Higuchi	0.951±0.027	73.42±7.22	2.79±0.55
Korsmeyer-Peppas	0.994±0.002	48.86±4.80	4.83±0.46
Hixson-Crowell	0.983±0.009	60.26±7.91	3.88±0.75
Hopfenberg	0.988±0.007	57.81±6.22	4.09±0.60
Baker-Lonsdale	0.910±0.033	81.67±4.91	2.10±0.35
Weibull	0.985±0.009	59.94±6.38	3.91±0.55

Table 5. Dissolution parameters of aspirin floating tablets by Korsmeyer-Peppas

Parameter	Mean ± SD
k _{KP}	1.175 ± 0.562
n	0.712 ± 0.091




Figure 1. Drug release of aspirin floating tablets




Figure 2. Observed data of drug release of aspirin floating tablets in dissolution studies, presented together with the model prediction using zero order, first order, Higuchi, Hixson-Crowell, Hopfenberg, Baker-Lonsdale, Korsmeyer-Peppas, and Weibull




Figure 3. Goodness of fit (GOF) evaluations based on correlation of residual versus time indicating that Korsmeyer-Peppas models do not form a specific pattern

**MAJALAH FARMASI INDONESIA
(INDONESIAN JOURNAL OF PHARMACY)**

ISSN : 0126-1037
STT. No. 1652/SK/DITJEN PPG/STT/1990
(Formerly SIT No. 1404/1972)

Email : mfi@ugm.ac.id; mfi.ugm@gmail.com
Website : <http://mfi.farmasi.ugm.ac.id>



MANUSCRIPT EVALUATION FORM

Date : 25-02-2015

Manuscript number : MFI-2015-02

Deadline : 11-03-2015

Title of Manuscript : In Vitro Release Modeling of Aspirin
Floating Tablets Using DDSolver

Please read the information below before reviewing the manuscript

1. Please return the reviewed manuscript within THREE WEEKS after the date of this manuscript evaluation form.
2. If reviewers can not review the manuscript due to DIFFERENT SCIENTIFIC FIELDS or the DEADLINE, please return the manuscript without delay.
3. The reviewers should give reviews and comments by considering the scientific benefit including research originality, adequacy and accuracy of research methodologies and techniques used, adequacy and accuracy of data analysis, comprehensiveness of literature review, and the readability and presentation of the article.
4. Please provide REVIEWS and COMMENTS to all items of the manuscript. The reviewers should provide DETAILED SUGGESTIONS for revision.

Please, express your opinion in relation to the manuscript (answer yes, no or partially):

1. Does the paper contain new information? **Y**
2. Does the title of the manuscript cover its content? **Y**
3. Is the summary appropriate? **Y**
4. Is the aim of the work clearly established? **Y**
5. Are the methods adequate? **Y**
6. Are the results reliable? **Y**
7. Are discussion and conclusions well documented and scientifically coherent? **Y**

Comments and suggestions for improvement:

EVALUATION REPORT (COMMENT SHEET)

Please provide some general comments to the authors in all items, and indicate the reason for your recommendation in all item. Please refer to the information no. 3 in page 1.

Please mark with (-) if there is no comment on each item.

Reviewer could use ADDITIONAL PAGES if column space provided is insufficient.

1.	Title The title describe the content of manuscript
2.	Abstract Fluidity is the tablet mass parameter, not tablet parameter, please this statement was corrected
3.	Introduction Attention for deleted word (typed in blue) and addition/correction words (typed in red)
4.	Methodology At analysis part: Attention for deleted word (typed in blue) and addition/correction words (typed in red)
5.	Result and Discussion Add some explanation here (type in red word)
6.	Conclusion -
7.	Acknowledgment (if any)

Recommendations

- | | | | |
|------------------------------|-------------------------------------|--|--------------------------|
| - Accept the manuscript | <input type="checkbox"/> | - Accept with major and extensive revision | <input type="checkbox"/> |
| - Accept with minor revision | <input checked="" type="checkbox"/> | - Reject the manuscript | <input type="checkbox"/> |

Please return the review and manuscript to :

Editorial

Majalah Farmasi Indonesia

(Indonesian Journal of Pharmacy)

Faculty of Pharmacy, Universitas Gadjah Mada,

Sekip Utara, Yogyakarta Indonesia, 55281. Telp: 62-274-6645911, 62-274-6492566

Fax : 62-274-543120

Indonesian J. Pharm. is indexed in Google Scholar, Index Copernicus, GetCITED, Genamics Journal Seek New Jersey USA, Geneva Foundation for Medical Education and Research (GFMER), Pharmaceutical Sciences Open Access Resources (PSOAR), International Pharmaceutical Abstracts (IPA), Inherent Digital Library UGM, WorldCat, Socolar, SCIRUS Elsevier, Indonesian Scientific Journal Database (ISJD), and being processed to be indexed in Academic Journal Database, DOAJ, CABI, ISI Thompson, PUBMED, Biological abstract, EBSCO Publishing Database. The journal is under planning to indexed with Pro-Quest, Scopus, Open J Gate.



Temukan pesan, dokumen, foto, atau orang



Awal

Tulis

← Kembali ↩️ ↶️ ➡️ 📁 Arsipkan 📁 Pindahkan 🗑️ Hapus 🔒

📧 7 📄 ?



gus_ump@... 999+
agussiswanto1... 41

• Acknowledgement Manuscript Submission MS MFI-2015-02 Yahoo/Email M... ★

...
Email Masuk 999+
Belum Dibaca
Berbintang
Draft 154
Ter kirim
Arsip
Spam
Sampah
^ Lebih sedikit

• **mfi@ugm.ac.id** 📧 Sen, 12 Jan 2015 jam 09.36 ★
Kepada: gus_ump@yahoo.com

Dear Mr. Agus Siswanto

This message is to acknowledge the receipt of the manuscript you submitted via email to the Indonesian Journal of Pharmacy. Your manuscript ID:MFI-2015-02. Entitle : "In Vitro Release Modeling of Aspirin Floating Tablets Using DDSolver"
Your manuscript has been received. Your paper will be reviewed by three or more reviewers assigned by the editorial board and final decision made by the editor will be informed by this email. You can monitor the review process for your paper by emailing us on the "Status of my manuscript No. MFI-2015-02".

Thank you very much for submitting your manuscript to the Indonesian Journal of Pharmacy.

Sincerely yours,

Prof. Dr. Abdul Rohman, M.Si., Apt
Associate editor
Indonesian Journal of Pharmacy

Info UGM: Sampaikan saran dan keluhan terhadap layanan UGM melalui sistem informasi aspirasi <http://aspirasi.ugm.ac.id>

↩️ ↶️ ➡️ ⋮

Balas, Balas ke Semua atau Teruskan



mfi@ugm.ac.id 🔍

mfi@ugm.ac.id
+ Tambah ke kontak

Asuransi Mobil Dijamin Terher

All Risk Mulai
~~Rp 3.050.000/tahun~~
Rp 1.982.500/tahun

- 500+ Bengkel Rekanan
- Usta Mobil s.d 15 Tahun
- Klaim Mudah Semua Kerusakan

CEK SEKARANG



Asuransi Mobil Dijamin Terher

All Risk Mulai
~~Rp 3.050.000/tahun~~
Rp 1.982.500/tahun

- 500+ Bengkel Rekanan
- Usta Mobil s.d 15 Tahun
- Klaim Mudah Semua Kerusakan

CEK SEKARANG





Temukan pesan, dokumen, foto, atau orang



Awal

Tulis

← Kembali ↶ ↷ ➡ 📁 Arsipkan 📁 Pindahkan 🗑️ Hapus 🔒

📧 📅 📄 ?



gus_ump@... 999+

agussiswanto1... 41

Email Masuk 999+

Belum Dibaca

Berbintang

Draft 154

Terkirim

Arsip

Spam

Sampah

^ Lebih sedikit

Tampilan Sembunyikan

📷 Foto

📄 Dokumen

📧 Langganan

✂️ Promo

Folder Sembunyikan

+ Folder Baru

• Bls: Review manuscript IJP-2015-2 Yahoo/Terkirim ★

AS • **agus siswanto** <gus_um> 📄 🔒 Rab, 18 Mar 2015 jam 09.20 ★
Kepada:
mfi.ugm@gmail.com

> Tampilkan pesan asli



In Vitro Rele... .doc
2.2MB

↶ ↷ ➡ ⋮

Balas, Balas ke Semua atau Teruskan

Kirim 📎 📄 📄 📄 📄 🔗 B I 🗑️

Asuransi Mobil Dijamin Terher

All Risk Mulai
~~Rp 3.050.000/tahun~~
Rp 1.982.500/tahun

- 500+ Bengkel Rekanan
- Usia Mobil s.d 15 Tahun
- Klaim Mudah Semua Kerusakan

CEK SEKARANG



Asuransi Mobil Dijamin Terher

All Risk Mulai
~~Rp 3.050.000/tahun~~
Rp 1.982.500/tahun

- 500+ Bengkel Rekanan
- Usia Mobil s.d 15 Tahun
- Klaim Mudah Semua Kerusakan

CEK SEKARANG



IN VITRO RELEASE MODELING OF ASPIRIN FLOATING TABLETS USING DDSOLVER

Agus Siswanto^{1,2}, Achmad Fudholi^{1*}, Akhmad Kharis Nugroho¹, Sudibyo Martono¹

¹Faculty of Pharmacy,
Gadjah Mada University,
Sekip Utara Jogjakarta
55281, Indonesia

²Faculty of Pharmacy,
Muhammadiyah University
of Purwokerto, Jl. Raya
Dukuh Waluh Purwokerto
53142, Indonesia

Submitted: 14-01-2015

Revised: 10-02-2015

Accepted: 25-03-2015

*Corresponding author
Achmad Fudholi

Email:
fudholiugm@yahoo.com

ABSTRACT

Aspirin has low solubility in water therefore, dissolution is a rate limiting step for absorption. Floating tablet formulation is designed to improve the bioavailability of aspirin. The objective of this study was to determine in vitro dissolution study of aspirin floating tablet release kinetics model. The floating tablets were prepared by a direct compression method using Methocel K4M CR, NaHCO₃, Ethocel, Aerosil, and dicalcium phosphate anhydrous as excipients. Tablets were evaluated by different parameters such as physicochemical properties, floating lag time ($F_{lag\ time}$), total floating time, and dissolution. The result showed that the tablet mass has good flow properties of 13.54 g/sec. Aspirin floating tablets had a weight uniformity (CV=1.45%), good hardness (6.42kg), and low friability (0.158%). The tablet has a short $F_{lag\ time}$ of 25.16 sec and long floating time of 8 hours. Dissolution data were evaluated using DDSolver conducted by (1) Statistical parameters: $R^2_{adjusted}$, AIC, MSC; (2) Visual goodness of fit (GOF). The results showed that aspirin floating tablets release kinetics followed the Korsmeyer-Peppas model. Aspirin release occurs through the mechanism of anomalous transport which combines Fickian diffusion and polymer relaxation.

Key words: aspirin floating tablet, DDSolver, modeling of drug release

INTRODUCTION

Aspirin is used as an antiplatelet for stroke prophylaxis (Laurer, 2002; Awtry and Loscalzo, 2000; Patrono and Rocca, 2008). Aspirin prevents blood from clotting by inhibiting cyclooxygenase, the enzyme responsible for the formation of thromboxane A₂, which is a mediator of platelet aggregating (Kannan *et al.*, 2010; Awtry and Loscalzo, 2000).

Aspirin has a low bioavailability due to first-pass effect and hydrolysis to salicylate metabolism in the gut wall (Sweetman, 2009). Aspirin is rapidly absorbed in the upper gastrointestinal tract, especially in the portion of the small intestine (Awtry and Loscalzo, 2000; Sweetman, 2009). Therefore, formulation of floating drug delivery system (floating on the gastric fluid) are designed to improve the bioavailability of aspirin (Suratri, 2008).

Aspirin has low solubility in water (1:300) (Moffat *et al.*, 2011; Sweetman, 2009) therefore, dissolution is one of the rate limiting step in their absorption and bioavailability (Gordon *et al.*, 1994; Wells, 2002). Information

on the mechanisms and kinetics of dissolution is important to estimate the absorption of aspirin floating tablet. Several mathematical models were developed to analyze dissolution data such as the zero-order, first-order, Higuchi, Weibull, Korsmeyer-Peppas, Hixson-Crowell, Bakar-Lonsdale, and Hopfenberg. The nonlinear fitting of dissolution data can be performed using a professional statistical programs such as Micro-Math Scientist, GraphPad Prism, SigmaPlot, or SYSTAT. However, these programs require the user to define the equations manually and need the initial value for each parameter (Zhang *et al.*, 2010). DDSolver is a new software developed for the kinetic analysis of dissolution data (Murtaza *et al.*, 2012) with a non-linear regression approach. DDSolver is a menu-driven add-in program for Microsoft Excel written in Visual Basic for Applications. Calculation using Excel offers a number of advantages over other software. The most attractive of advantages is the ease of use. Most scientists are already familiar with Excel because of its wide availability and high

flexibility (Zhang *et al.*, 2010). This study was aimed to describe the dissolution data of aspirin floating tablet using DDSolver.

MATERIAL AND METHODS

Aspirin was obtained from Jilin Pharmaceutical Co. (Jilin, China), sodium bicarbonate from Tosoh Corporation (Tokyo, Japan), dicalcium phosphate anhydrous from Budenheim (Budenheim, Germany), Methocel K4M CR and Ethocel Premium Standard 20 from Colorcon Asia Pvt. Ltd. Hydrochloric acid and sodium chloride (analytical grade) were purchased from Merck (Darmstadt, Germany). All other chemicals used were of pharmaceutical grade.

Preparation of floating tablet

Tablets were prepared by direct compression method based on formulation in Table I. Aspirin, Methocel K4M CR, sodium bicarbonate, Ethocel, dicalcium phosphate anhydrous, and aerosil were mixed in a cube mixer at 30rpm for 3min. The tablet compression process used single punch tablet machine (Korsch, Germany). The weight of tablets were arranged at 200mg and the hardness of tablet were controlled at 6-8kg (Monsanto hardness tester).

Table I. Formulation of aspirin floating tablet

Ingredient	Amount (mg) / tablet
Aspirin	80.00
Methocel K4M CR	33.87
NaHCO ₃	30.37
Ethocel	35.76
Aerosil	2.00
Dicalcium phosphate anhydrous	18.00
Total weight (mg)	200.0

Physical properties of tablet

Weight variation: Twenty tablets were randomly selected from the batch and weighed individually to check for weight variation. Hardness test: Monsanto hardness tester was used to determine the tablet hardness. Friability: About 20 tablets were selected from each batch and evaluated for

friability using an abrasive tester (ERWEKA, Germany).

Floating lag time (F_{lag time})

Tablet was immersed in beaker containing 100 mL of simulated gastric fluid (SGF) without pepsin pH 1.2. Time required for the tablet to rise to the surface and floating was determined as floating lag time (Patel *et al.*, 2007a).

In vitro dissolution studies

The in vitro dissolution study was determined using Erweka DT 600 USP apparatus II (paddle method), using 900mL SGF without pepsin pH 1.2 as dissolution medium at 37±0.2°C and 60rpm (USP XXXII, 2008; Gopalakrishnan and Chentilnathan, 2011). Aliquots of 5mL was taken out at intervals of 15, 30, 45, 60, 120, 180, 240, 300, 360, 420, and 480min. Exactly 5mL of fresh SGF was added to the dissolution vessel after each withdrawal to maintain a constant volume. The samples were analyzed by UV-Vis spectrophotometer (Genesys 10 Thermo) at 280nm.

Determination of drug content

Twenty tablets were taken randomly from the batch, weighed, and powdered. A 200mg of powder equivalent to 80mg of aspirin was put into a 100mL flask. Two mL of absolute ethanol and simulated gastric fluid (without pepsin) were added up to the mark. The solution was centrifuged for 60min, then filtered with filter paper. The drug content was analyzed at the wavelength of aspirin (278nm) and salicylic acid (304nm) using a UV-Vis spectrophotometer. Drug content was then calculated with a simultaneous spectrophotometric method.

Data analysis

Physicochemical and floating properties were evaluated based on the applicable requirements. The dissolution data modeling was calculated using DDSolver. Evaluation of release kinetics model was conducted by (1) Statistical parameters: R²_{adjusted}, Akaike Information Criterion (AIC), Model Selection Criterion (MSC); and (2) Visual goodness of fit (GOF).

**RESULTS AND DISCUSSION****The physicochemical properties of aspirin floating tablets**

Aspirin floating tablets have good physicochemical properties (Table II). The tablets have a uniform weight with CV=1.45 % so the active substance in each tablet was uniform (Davies, 2009). Tablets have good hardness of 6.42kg (Gordon *et al.*, 1990) and low friability (<1%) therefore, it withstand to mechanical shocks and not easily destroyed. The content of aspirin in floating tablets within the required USP XXXII (2008) range of 95-105%.

Table II. Physicochemical properties of aspirin floating tablets

Parameters	Mean \pm SD
Weight of tablet (mg)	201.1 \pm 0.6
Friability (%)	0.16
Hardness (kg)	6.42 \pm 0.58
F _{lag time} (sec)	25.16 \pm 4.60
Total floating time (hours)	>8
Drug content (aspirin) (%)	99.55 \pm 1.99
Drug content (salysilic acid) (%)	1.24 \pm 0.09

**In vitro buoyancy of aspirin floating tablets**

One parameter of good floating tablet quality is the tablet's ability to float in the medium indicated by F_{lag time} and floating time. The duration of floating indicates the time of tablets float in the dissolution medium. Aspirin floating tablets was able to float during dissolution test, intact and not eroded. This indicates that the floating tablet of aspirin can release during dissolution testing. The composition of excipients used in the formulation was able to maintain the floating ability and the integrity of the tablets. Ethocel, Aerosil, and dicalcium phosphate anhydrous are hydrophobic excipients that play a role in the maintaining of integrity of the tablets in the dissolution medium.

F_{lag time} indicates the speed of tablet to float in the dissolution medium. F_{lag time} test (Table II) showed that aspirin floating tablets within the requirements of less than 180 seconds (Patel *et al.*, 2007a). The sooner the tablet floats, the better the floating quality.

Buoyancy was determined by the density of tablet. If the density is less than 1, the tablets will float properly (Patel *et al.*, 2007a; Dave *et al.*, 2004). Aspirin floating tablet buoyancy is determined by the composition of Methocel K4M CR, NaHCO₃, and Ethocel as excipients. When NaHCO₃ reacts with SGF, the CO₂ gas was produced. Gas trapped in a hydrophilic gel Methocel K4M CR thereby, decreasing the density of the tablets (Patel *et al.*, 2007b). Ethocel, which is hydrophobic, will maintain the integrity of tablet to float in a long time. Therefore, the combination of these three excipients produced a floating tablet with short F_{lag-time} and long floating duration.

In vitro dissolution profiles of aspirin floating tablets

Aspirin floating tablet dissolution test was conducted according to USP XXXII (2008). Aspirin tablets were designed to float in the stomach for a long time. Drug release will occur in the gastric; therefore, the SGF (without pepsin) pH 1.2 was used as dissolution test medium. The dissolution profile was made by plotting the percentage of drug dissolved versus time (Figure 1). The amount of aspirin dissolved in the 60 minutes is 22.73% within USP XXXII (2008) range of 20-55%. The amount of aspirin released after 8h was 84.85%.

Dissolution Study of Aspirin Floating Tablet Release using DDSolver

In vitro drug release data were fitted to kinetics models such as the zero-order, first-order, Higuchi, Weibull, Korsmeyer-Peppas, Hixson-Crowell, **Bakar-Lonsdale**, and Hopfenberg using DDSolver. The input of dissolution data into DDSolver include: (1) percentage of drug dissolved, (2) dissolution time (minutes). Modeling dissolution by DDSolver were estimated by nonlinear regression in accordance with the equation in table III.

Analysis model of visual goodness of fit (GOF) conducted in two methods: (1) the correlation of Q_o (amount of aspirin dissolved in the experimental results) versus Q_c (the predicted amount of aspirin dissolved in the medium) (Figure 2); (2) the correlation of residuals versus time (Figure 3) (Nugroho *et al.*, 2014).

Table III. Models available in **DDsolver** for fitting drug release data

Model	Equation
Zero order	$F=k_0 t$
First order	$F=100 [1-\text{Exp}(-k_1 t)]$
Higuchi	$F=k_H t^{0.5}$
Hixson-Crowell	$F=100 [1-(1-k_{HC} t)^3]$
Hopfenberg	$F=100 [1-(1-k_{HB} t)^n]$
Weibull	$F=100 \{1-\text{Exp}[-(t^\beta)/\alpha]\}$
Korsmeyer-Peppas	$F=k_{KP} t^n$
Baker-Lonsdale	$3/2 [1-(1-F/100)^{2/3}]-F/100=k_{BL} t$

Note: F = drug released (%); t = time (minutes)




Figure 1. Drug release of aspirin floating tablets

Correlation analysis of Q_0 versus Q_c in figure 2 showed that the Korsmeyer-Peppas model was the most appropriate model to explain the phenomenon of aspirin dissolution in floating tablet. Kormeyer-Peppas models provide the less deviation between Q_0 and Q_c , so it appears in figure 2 that the distribution of dissolution data (Q_0) beyond the curve of predicted dissolution data (Q_c). Modeling kinetics of dissolution in zero-order, first-order, Higuchi, Weibull, Hixson-Crowell, **Bakar-Lonsdale**, and Hopfenberg produced the more deviation of Q_0 and Q_c . This is reinforced by the residual correlation analysis (Q_0-Q_c) versus time indicates that Korsmeyer-Peppas models do not form a specific pattern in figure 3.

DDSolver provides a number of statistical criteria to evaluate the dissolution model included the correlation coefficient ($R_{\text{obs-pre}}$), the coefficient of determination (R_{sqr} , R^2 , or COD), the adjusted coefficient of determination ($R_{\text{sqr_adj}}$ or R^2_{adjusted}), mean square error (MSE), standard deviation residuals (MSE_{root} or $Sy.x$), SS, WSS, Akaike Information Criterion (AIC), and Model Selection Criterion (MSC). Among these parameters, R^2_{adjusted} , AIC, MSC are the most popular and widely used in the identification of dissolution data modeling (Zhang *et al.*, 2010).

R^2_{adjusted} is the most appropriate parameter for comparing dissolution models with different parameters compared to R^2 .




Figure 2. Observed data of drug release of aspirin floating tablets in dissolution studies, presented together with the model prediction using zero order, first order, Higuchi, Hixson-Crowell, Hopfenberg, Baker-Lonsdale, Korsmeyer-Peppas, and Weibull.




Figure 3. Goodness of fit (GOF) evaluations based on correlation of residual versus time indicating that Korsmeyer-Peppas models do not form a specific pattern

Table IV. Statistical parameters of models to describe release of aspirin floating tablets

Model	Statistical parameters of models (mean±SD)		
	R ² _{adjusted}	AIC	MSC
Zero order	0.938±0.037	75.38±8.65	2.62±0.77
First order	0.982±0.010	61.35±5.98	3.79±0.51
Higuchi	0.951±0.027	73.42±7.22	2.79±0.55
Korsmeyer-Peppas	0.994±0.002	48.86±4.80	4.83±0.46
Hixson-Crowell	0.983±0.009	60.26±7.91	3.88±0.75
Hopfenberg	0.988±0.007	57.81±6.22	4.09±0.60
Baker-Lonsdale	0.910±0.033	81.67±4.91	2.10±0.35
Weibull	0.985±0.009	59.94±6.38	3.91±0.55

Table V. Dissolution parameters of aspirin floating tablets by Korsmeyer-Peppas

Parameter	Mean ± SD
k _{KP}	1.175 ± 0.562
n	0.712 ± 0.091

R²_{adjusted} value will decrease with the increasing of dissolution model parameters. The best model is the model with the highest value of R²_{adjusted} (Zhang *et al.*, 2010). Based on R²_{adjusted} in table IV, Korsmeyer-Peppas is the most appropriate model for this study.

AIC parameters have been used for more than 35 years in the dissolution data analysis because it is simple and easy to use (Zhang *et al.*, 2010). The model with the smallest AIC value is the most precise model (Motulsky and Christopoulos, 2003). The MSC is a modified reciprocal of the AIC and has been normalized so that it is independent of the scaling of the data points. If we compared several models, the model with the highest value of the MSC is the best model. In general, the model was appropriate when the MSC value > 2 or 3 (Zhang *et al.*, 2010). Dissolution model parameters in table IV showed that the Korsmeyer-Peppas models is the best model to explain the phenomenon of aspirin floating tablet dissolution.

Dissolution mechanism of aspirin floating tablet by Korsmeyer-Peppas

Korsmeyer-Peppas dissolution models explain the exponential relationship of dissolved drug fraction versus time.

$$F = k_{KP} t^n$$

F = fraction of drug dissolved at time t, a = drug dissolution constant, t = dissolution time (minutes), and n = dissolution exponent (Costa and Lobo, 2001).

Dissolution exponent (n) can be used to explain the mechanisms involved in the dissolution (Kalam *et al.*, 2007). The analysis result of aspirin floating tablet dissolution with Korsmeyer-Peppas gave n value of 0.712 (Table V). For cylindrical dosage form (tablets) then 0.45 < n < 1 indicates that dissolution occurs through the mechanism of anomalous transport (Costa and Lobo, 2001; Colombo *et al.*, 2000). The release of aspirin from floating tablet occurs through the combination of Fickian diffusion and polymer relaxation. The dissolution of aspirin was influenced by the excipients in the tablet. Methocel K4M CR is a hydrophilic polymer HPMC. When it contacted with SGF, Methocel K4M CR will expand to form a thick gel. NaHCO₃ reacts with dissolution medium (SGF) produces CO₂ which form pores in the polymer gel Methocel K4M CR. Ethocel, which is hydrophobic, will maintain the integrity of the tablets to withdraw the massive erosion. Therefore, the release of aspirin occurs through Fickian diffusion pass through hydrophilic gel. The dissolution of aspirin was also determined by the relaxation ability of Methocel K4M CR and Ethocel in the medium.

CONCLUSION

Based on the results it can be concluded that aspirin floating tablets fulfilled the requirements of good quality tablet. The dissolution data evaluated with DDSolver showed that the kinetics release of floating tablets followed the Korsmeyer-Peppas model through anomalous transport mechanism which combines Fickian diffusion and polymer relaxation.

ACKNOWLEDGMNET

The authors express their sincere thanks to Faculty of Pharmacy, Universitas Gadjah Mada, and Faculty of Pharmacy, Muhammadiyah University of Purwokerto

REFERENCES

- Awtry EH., Loscalzo J., 2000, Aspirin, *Circulation*, **101**, 1206-1218
- Costa P., Lobo JMS., 2001, Review Modeling and comparison of dissolution profiles, *Eur. J. Pharm. Sci.*, **13**, 123-133
- Colombo P., Santi P., Bettini R., Brazel CS., Peppas NA., 2000, Drug Release from Swelling-Controlled System, in Wise, D.L., Handbook of Pharmaceutical Controlled Technology, *Marcel Dekker Inc., New York*. pp 183-205
- Dave BS., Amin AF., Patel MM., 2004, Gastroretentive Drug Delivery System of Ranitidine Hydrochloride: Formulation and In Vitro Evaluation, *AAPS PharmSciTech*, **5(2)**, 1-6
- Davies P., 2009, Oral Solid Dosage Form, In Gibson M., Pharmaceutical Preformulation and Formulation, *Informa healthcare, New York*. pp 389
- Fudholi A., 1983, Methodology Formulation in Direct Compression, *Medika*, **7(9)**, 586-593
- Gopalakrishnan, S., Chenthilnathan, A., 2011, Floating Drug Delivery Systems: A Review, *J. Pharm. Sci. Tec.*, **3(2)**, 548-554.
- Gordon RE., Rosanke TW., Fonner DE., Anderson NR., Banker GS., 1990, Granulation Technology and Tablet Characterization, in Lieberman, H.A., Lachman, L., Schwartz, J.B., Pharmaceutical Dosage Form: Tablet, 2nd Ed., Vol 2, *Marcel Dekker Inc.*, New York. pp 327-332
- Gordon MS., Ellis DJ., Molony B., 1994, In Vitro Dissolution Versus In Vivo Evaluation of Four Different Aspirin Products, *Drug Dev. Ind. Pharm.*, **20(10)**, 1711-1723
- Kannan S., Manivannan R., Balasubramaniam A., Kumar NS., 2010, Formulation and Evaluation of Aspirin Delayed Release Tablet, *IJCP*, 01, accessed 12.02.12, <<http://www.pharmacie-globale.info>>
- Kalam MA., Humayun M., Parvez N., Yadaz S., Garg A., Amin S., Sultana Y., Ali A., 2007, Release Kinetics of Modified Pharmaceutical Dosage Forms: A Review, *J. Pharm. Sciences*, **1**, 30-35
- Lauer MS., 2002, Aspirin For Primary Prevention of Coronary Events, *N Engl J Med*, **346(19)**, 1468-1474
- Moffat AC., Osselton MD., Widdop B., Watts J., 2011, Clarke's Analysis of Drugs and Poisons in Pharmaceuticals, Body Fluids and Postmortem Material, fourth edition, Pharmaceutical Press, London-Chicago. pp 924-926
- Motulsky HJ., Christopoulos A., 2003, Fitting Models To Biological Data Using Linear and Nonlinear Regression. A Practical Guide To Curve Fitting, GraphPad Software, Inc., San Diego CA. pp 143-148
- Murtaza G., Ahmad M., Khan SA., Hussain I., 2012, Evaluation of Cefixime-Loaded Chitosan Microspheres: Analysis of Dissolution Data Using DDSolver, *Dissolution Technologies*, 13-19
- Nugroho AK., Binnarjo A., Hakim AR., Ermawati Y., 2014, Compartmental Modeling Approach of Losartan Transdermal Transport In Vitro, *Indonesian J. Pharm.*, Vol. 25 No. 1, 31-38
- Patel DM., Patel NM., Pandya NN., Jogani, PD., 2007a, Gastroretentive Drug Delivery System of Carbamazepine: Formulation Optimization Using Simplex Lattice Design: A Technical Note, *AAPS PharmSciTech*, **8(1)**, E1:E5, accessed 24.10.11, <<http://www.aapspharmscitech.org>>
- Patel VF., Patel NM., 2007b, Statistical Evaluation of Influence of Viscosity and Content of Polymer on Dipyrindamole



- Release From Floating Matrix Tablets: A Technical Note, *AAPS PharmSciTech*, **8(3)**, E2-E5, accessed 24.10.11, <<http://www.aapspharmscitech.org>>
- Patrono C., Rocca B., 2008, Aspirin: Promise and Resistance in the New Millennium, *Arterioscler Thromb Vasc Biol*, **28**, s25-s32, accessed 16.11.11, <<http://atvb.ahajournals.org>>.
- Suratri A., 2008, Floating Drug Delivery Systems To Increase Gastric Retention of Drugs: A Review, *RJPT*, **1(4)**, accessed 02.04.11, <<http://www.rjptonline.org/>>.
- Sweetman SC., 2009, Martindale The Complete Drug Reference, Thirty-sixth edition, *Pharmaceutical Press*, London. pp 20-25
- USP, 2008, United States Pharmacopeial XXXII, accessed 17.02.12, <<http://www.pharmacopeia.cn/>>.
- Wells J., 2002, Pharmaceutical preformulation: The Physicochemical Properties of Drug Substances, in Aulton, M.E., *Pharmaceutics: The Science of Dosage Form Design*, Ed. II, *Churchill Livingstone*, Edinburg. pp 113-138
- Zhang Y., Huo M., Zhou J., Zou A., Li W., Yao C., Xie S., 2010, DDSolver: An Add-In Program for Modeling and Comparison of Drug Dissolution Profiles, *The AAPS Journal*, **Vol. 12 No. 3**, 263-271

Galley Proof



Temukan pesan, dokumen, foto, atau orang



Awal

Tulis

Kembali Arsipkan Pindahkan Hapus

7 ?



gus_ump@... 999+
agussiswanto1... 41

Email Masuk 999+
Belum Dibaca
Berbintang
Draft 154
Ter kirim
Arsip
Spam
Sampah
^ Lebih sedikit

Tampilan Sembunyikan

Foto
Dokumen
Langganan
Promo

Folder Sembunyikan

+ Folder Baru

Revisi gallery proof (agus siswanto) Yahoo/Email M...
In vitro release modeling... 2

AS • agus siswanto <gus_ump@gmail.com> Rab, 29 Apr 2015 jam 08.55
Kepada: mfi.ugm@gmail.com
Cc: siswanto.kpp@gmail.com



Galley Proofpdf
906.3kB

Navigation icons: back, forward, refresh, etc.

M • mfi ugm <mfi.ugm@gmail.com> Rab, 29 Apr 2015 jam 10.06
Kepada: gus_ump@yahoo.com

ini mas
> Tampilkan pesan asli



5. 94-102 ag....doc
433kB

Asuransi Mobil Dijamin Terher

All Risk Mulai
~~Rp 3.050.000/tahun~~
Rp 1.982.500/tahun

- 500+ Bengkel Rekanan
- Usia Mobil s.d 15 Tahun
- Klaim Mudah Semua Kerusakan

CEK SEKARANG

Analisis keuangan pribadi

Dapatkan akses strategi tradin yang dikembangkan oleh trade expert.



Temukan pesan, dokumen, foto, atau orang



Awal

Tulis

← Kembali ↩️ ↶️ ➡️ 📁 Arsipkan 📁 Pindahkan 🗑️ Hapus 🔒

📧 7 📄 ?



gus_ump@... 999+

agussiswanto1... 41

Email Masuk 999+

Belum Dibaca

Berbintang

Draft 154

Terkirim

Arsip

Spam

Sampah

^ Lebih sedikit

Tampilan Sembunyikan

📷 Foto

📄 Dokumen

📧 Langganan

✂️ Promo

Folder Sembunyikan

+ Folder Baru

revisi artikel

Yahoo/Terkirim ★



• agus siswanto <gus_ump@...>
 Kepada: mfi.ugm@gmail.com
 Cc: siswanto.kpp@gmail.com

Rab, 29 Apr 2015 jam 11.21 ★



5. 94-102 ag... .doc
 428.5kB

↩️ ↶️ ➡️ ⋮

Balas, Balas ke Semua atau Teruskan

Kirim 📎 📄 📧 😊 🔗 B I 🗑️

Asuransi Mobil Dijamin Terher

All Risk Mulai ~~Rp 3.050.000/tahun~~
Rp 1.982.500/tahun

- 500+ Bengkel Rekanan
- Usia Mobil s.d 15 Tahun
- Klaim Mudah Semua Kerusakan

CEK SEKARANG



Asuransi Mobil Dijamin Terher

All Risk Mulai ~~Rp 3.050.000/tahun~~
Rp 1.982.500/tahun

- 500+ Bengkel Rekanan
- Usia Mobil s.d 15 Tahun
- Klaim Mudah Semua Kerusakan

CEK SEKARANG

