

ANTIBACTERIAL, IN VITRO CYTOTOXIC, AND ANTIOXIDANT ACTIVITIES OF ELECTROLYZED OXIDIZING/REDUCING WATER

By asmiyenti djaliasrin djali

ANTIBACTERIAL, INVITROCYTOTOXIC, AND ANTIOXIDANT ACTIVITIES OF ELECTROLYZED OXIDIZING/REDUCING WATER

ASMIYENTI DJALIASRIN DJALIL^{1*}, ELZA SUNDHANI¹, RETNO WAHYUNINGRUM¹, DWI HARTANTI¹, NONI LESTIOWATI¹, PRAYOGO PANGESTU¹, RIDWAN SETIAJI¹, SUWANDRI SUWANDRI²

¹Department of Pharmacy, Faculty of Pharmacy, Universitas Muhammadiyah Purwokerto, Jl. Raya Dukuwaluh, PO. Box 202, Purwokerto 53182, Indonesia. ²Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Jl. Dr. Soeparno, Purwokerto 53123, Indonesia. Email: asmientidjaliasrindjalil@ump.ac.id

Received: 10 November 2018, Revised and Accepted: 06 December 2018

ABSTRACT

Objective: Electrolyzed oxidizing/reducing water is popular as health beneficial water in Indonesia. In this study, we examined the level of antibacterial, anticancer, and antioxidant activity of the electrolyzed water.

Methods: The efficacy of electrolyzed water produced by Enagic® at six level pH (2.5, 6.0, 7.0, 8.5, 9.0, and 9.5) was investigated. Antibacterial activity was evaluated using a macrodilution method. The anticancer activity was performed against human breast cancer (T47D) cell lines using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assay. Moreover, the antioxidant activity was determined using antioxidant model, 2,2-diphenyl-1-picrylhydrazyl radical scavenging activity.

Results: The results show that electrolyzed water exhibited antibacterial activity against *Propionibacterium acnes* and *Staphylococcus epidermidis*. Among six level pH, electrolyzed water at pH 2.5 showed the highest antibacterial activity. The *in vitro* cytotoxic activity of electrolyzed waters showed potential moderate cytotoxicity. The activity tends to be higher in alkaline electrolyzed water. However, the electrolyzed water showed free radical scavenging activity.

Conclusion: Electrolyzed water that marked in Indonesia has some potential health benefits. The activity is dependent on pH.

Keywords: Antibacterial activity, Antioxidant, Anticancer, Electrolyzed water, A healthy water.

© 2019 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ijap.2019.v11i5.T0043>

INTRODUCTION

Today, people are very concerned about consuming drinks that have some health beneficial effects, including the Indonesian community. The functional water can be obtained by irradiation of light, ultrasonication, treatment with a magnetic field, and electrolysis. Among these methods, electrolyzed water has been widely used for functional water.

Electrolyzed oxidizing/reducing water that has been developed in Japan is one option that has some health beneficial effects. This water is made through an ionization process by flowing electric current, which ultimately produces alkaline and acidic water. The alkaline water or electrolyzed reduced water (ERW) is rich in hydroxyl ions (OH⁻). On the other hand, the acidic water or electrolyzed oxidized water (EOW) is rich in hydrogen ions (H⁺). Its water has advantages because it has smaller molecular cluster compared to its raw material, making it more simply absorbed by the body. Electrolyzed water is made by passing the aqueous solution of NaCl or KCl solution on an electrolyzing machine. This machine consists of anodes and cathodes separated by a special diaphragm which separates the alkaline fraction at the cathode and the acid fraction at the anode [1]. The negatively charged particles and ions (Cl⁻ and OH⁻) move toward the anode where electrons are released, and hydrochloric acid (HCl), hypochlorous acid (HOCl), hypochlorite ion (OCl⁻), oxygen gas (O₂), and chlorine gas (Cl₂) are produced. However, positively charged ions and particles (H⁺ and Na⁺) move toward the cathode, where they obtain electrons, resulting in the generation of sodium hydroxide (NaOH) and hydrogen gas [2].

Previous studies have shown that EOW effectively kills *Escherichia coli*, *Salmonella enteritidis*, and *Listeria monocytogenes* [3,4]. The electrolyzed water shows bactericidal, virucidal, and fungicidal effects

in numerous areas such as medicine and dentistry as well as on fruits, vegetables, and agriculture [2,5]. ERW is known to suppress oxidative stress associated with the disease, such as diabetes [6], cancer [7-10], arteriosclerosis, and neurodegenerative diseases [11]. On the other hand, neutral electrolyzed water has been shown to be effective for the inactivation of human norovirus [12].

One of the most popular electrolyzed water in Indonesia is produced by Enagic. This water is available on the market at various pH values. The pH is 2.5, 6.0, 7.0, 8.5, 9.0, and 9.5 with different uses. This research was conducted to determine the electrolyzed water potential for health marked in Indonesia by conducting antibacterial, anticancer, and antioxidant activity tests.

MATERIALS AND METHODS

Materials

Water electrolyzed used for this study was purchased from local producers in Indonesia. The water was generated using a water ionization machine made by the Japanese company Enagic. Six different pH of electrolyzed waters were used in the present research. Human breast cancer T47D cells lines were provided from cancer and stem cells research center, Universitas Muhammadiyah Purwokerto. *Propionibacterium acnes* and *Staphylococcus epidermidis* were obtained from the American type culture collection (ATCC; Rockville, MD, USA) as well as the culture collection of the Laboratory Microbiology, Universitas Muhammadiyah Purwokerto. Fetal bovine serum (FBS), Dulbecco's modified eagles medium (DMEM), penicillin-streptomycin, trypsin-EDTA, and fungizone were purchased from Gibco (Invitrogen, USA). Phosphate-buffered saline (PBS) 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), 2,2-diphenyl-1-picrylhydrazyl

(DPPH), and nutrient agar (NA) were purchased from Sigma-Aldrich. Furthermore, sodium dodecyl sulfate (SDS) and HCl were obtained from Merck.

Antibacterial activity of electrolyzed water
Agar macrodilution was adopted for measuring the antibacterial activity of the electrolyzed water. Clindamycin (20 µg/mL) was used as the positive control; however, sterile distilled water was used as a negative control. Electrolyzed waters with different pH (2.5, 6.0, 7.0, 8.5, 9.0, and 9.5) were collected from Purwokerto, Indonesia. A 1 mL bacterial inoculum (1×10^8 colony-forming unit [CFU/mL]) was added to 9 mL aliquot of each electrolyzed water sample in 15 mL sterile conical tubes (5 tubes per samples). The tubes were incubated at 35°C for 6 min. The treatment electrolyzed waters were diluted serially and plated (100 µL per plate) on NA. Then, agar plates were incubated in an anaerobic environment at 37°C for 24 h. After incubation, the colony was counted by colony counter.

In vitro cytotoxic activity of electrolyzed water

Three different pH of electrolyzed waters (2.5, 6, and 9) were used in the experiment. T47D cells are regenerated in the medium when used *in vitro*. T47D cells were grown in DMEM growing media containing FBS 10% (v/v) and penicillin-streptomycin (100 IU/mL). The cells (10^4 cells/well) were planted into the 96 plates and incubated at 37°C for 24 h, with 5% CO₂. After incubation, the culture medium was removed and cells were washed using PBS. The cells were treated by electrolyzed water and then incubated again for 24 h. The concentration of each electrolyzed water was 100.0, 75.0, 50.0, 25.0, and 12.5% (v/v) in DMEM. Cell viability was determined by MTT assay. The cells were added with MTT-containing medium (0.5 mg/mL) for 4 h at 37°C, with 5% CO₂. Viable cells will react with MTT to generate purple colored formazan crystals. After 4 h of incubation, the reaction was stopped with 10% SDS in 0.1 M HCl solution. Then, the plate was incubated overnight at room temperature and protected from light. At the end of incubation, the plates were shaken, and cells absorbance was measured by ELISA reader (Bio-Rad) at λ 595 nm.

Antioxidant activity of electrolyzed water

Antioxidant activity of electrolyzed water was determined using the DPPH radical scavenging assay adopted by Brand-Williams *et al.* [13] with a minor modification. Briefly, the DPPH solution 0.004% (w/v) 2 mL was added to electrolyzed water in concentration from 6.25 to 100% (v/v). The mixture was allowed to stand at room temperature in the dark for 30 min. Then, absorbance is read at the maximum wavelength. After incubation, the absorbance was measured at 516 nm against a blank by spectrophotometer Ultraviolet (UV) Vis (Shimadzu UV-1201). The ability to scavenge the DPPH• was determined using the formula given by Brand-Williams *et al.* [13]. The activities were compared to L-ascorbic acid [14].

RESULTS AND DISCUSSION

In this research, six different commercially available pH of electrolyzed water were evaluated by pH meter (metrohm). These pH values are shown in Table 1. It can be observed that the pH values found for samples 2-5 are in close agreement with the labeled pH. Sample 1 presented pH value above the labeled pH, whereas sample 6 presented pH value under the labeled pH.

The highest difference of pH value is in the sample with a pH label 2.5. Based on data from chemical analysis and spectroscopic data, the main components of electrolyzed acidic water are Cl₂/HClO, which is in equilibrium according to the pH of the solution [4,15]. The Cl₂ gas is susceptible to decrease due to the storage; this will cause the pH of the water to increase.

Antibacterial activity of electrolyzed water

Antibacterial activity of electrolyzed water was investigated against skin poisoning bacteria, including *P. acnes* and *S. epidermidis*. The bacterial cells have been recognized as bacteria initiating inflammation in acne.

It has been claimed that electrolyzed water, especially marked in the strong acid water, can be used to treat infections caused by bacteria on the skin. Clindamycin antibiotics are used as a positive control. The results of electrolyzed water in killing *P. acnes* and *S. epidermidis* are reported in Table 2.

Unlike electrolyzed water at pH 6.0, 7.0, 8.5, 9.0, and 9.5, the results of the antibacterial activity test against *P. acnes* and *S. epidermidis* at pH 2.5 showed that there were no bacterial colonies growing (Fig. 1). This result is in line with previous studies which indicated that electrolyzed oxidizing water might be potential as a disinfectant [16]. This is possible because electrolyzed water at low pH contains a high concentration of HOCl. The concentration of HOCl is known to correlate with the ability of electrolyzed water to kill bacterial cells [15]. In water with a low pH can damage the outer membrane of bacterial cells so that HOCl can enter bacterial cells more efficiently [17]. HOCl is the most active chlorine compound, apparently killing microbial cells through inhibition of glucose oxidation by chlorine sulfide-oxidizing groups of certain enzymes that are important in carbohydrate metabolism [18].

Table 1: pH measurements of electrolyzed water

Sample	Labelled pH	Found pH*	pH differences
1	2.5	2.76±0.04	0.36
2	6.0	6.03±0.05	0.03
3	7.0	7.01±0.02	0.01
4	8.5	8.47±0.04	0.03
5	9.0	8.99±0.02	0.01
6	9.5	9.38±0.21	0.12

*Mean±SD, n= 3

Table 2: Viability of *P. acnes* and *S. epidermidis* after exposure to electrolyzed water measurements of electrolyzed water

Treatment	CFU/mL	
	<i>P. acnes</i>	<i>S. epidermidis</i>
Positive control	0±0	0±0
pH 2.5	0±0	0±0
pH 6.0	1420±148	398±69
pH 7.0	557±103	1927±661
pH 8.5	403±67	1116±117
pH 9.0	1168±175	228±21
pH 9.5	1503±122	24±3

*Mean±SD, n=5, *P. acnes*: *Propionibacterium acnes*, *S. epidermidis*: *Staphylococcus epidermidis*, CFU: Colony-forming unit




Fig. 1: Effect of electrolyzed water at pH 2.5 on the growth of *Propionibacterium acnes* (top) and *Staphylococcus epidermidis* (bottom) by agar macrodilution

Table 3: Antioxidant activity of electrolyzed water, using DPPH free radical scavenging method

Treatment	EC ₅₀ ±SD (%)	EC ₂₀ ±SD (%)
pH 2.5	227.4±9.7	102.5±8.0
pH 6.0	293.0±34.4	130.3±16.5
pH 7.0	607.4±28.7	219.6±13.1
pH 8.5	195.8±4.5	63.3±2.3
pH 9.0	177.4±1.6	63.2±4.0
pH 9.5	167.6±2.8	56.0±3.2
Ascorbic acid	9.3±0.4 (µg/mL)	3.4±0.3 (µg/mL)

DPPH: 2,2-diphenyl-1-picrylhydrazyl, EC₅₀: 50% efficient concentration

P. acnes grew well in neutral or slightly alkaline. This result is in line with previous studies [19]. We observed that *P. acnes* was still survived even at pH 9.5 of electrolyzed water. Furthermore, *S. epidermidis* grew much better in the pH of 7.0–8.5. However, minor changes in the pH in the pH 9.0–9.5 range can decrease the number of *S. epidermidis*. *S. epidermidis* did not survive at pH 9.5 of electrolyzed water. A substantial population reduction of *S. epidermidis* at pH 9.5 (24 CFU/mL) was observed after 24 h incubation. The reduction population of *S. epidermidis* is known to correlate with the high concentration of OCl⁻ in electrolyzed water at pH 9.0–9.5 [20].

In vitro cytotoxic activity of electrolyzed water

The *in vitro* cytotoxic activity of electrolyzed waters with different pH was evaluated against T47D breast cancer cell. T47D cell is a continuous cell line isolated from ductal breast tumor tissue. This cell expresses the p53 protein, which has a mutation. This causes p53 cannot bind to the response of elements to DNA. Thus, the ability of p53 to regulate cell cycle is reduced and can even remove [21]. The treatment of electrolyzed water samples for 24 h resulted in a decrease in percentage of T47D cell viability compared to cell control (Fig. 2). Under acidic, slightly acidic, and slightly alkaline conditions, electrolyzed waters had *in vitro* cytotoxic activity. The electrolyzed water at pH 2.5 had the lowest cytotoxicity on T47D cell lines, with an the half maximal inhibitory concentration (IC₅₀) of 59.4%, followed by electrolyzed water at pH 6 with an IC₅₀ value of 47.0%, and electrolyzed water at pH 9 with an IC₅₀ value of 46.6%.

Morphological alteration of T47D cell lines on exposure using electrolyzed water was demonstrated under an inverted microscope (Carl Zeiss) with a magnification of ×20 (Fig. 3). The cells showed the most noticeable effect after incubation to the electrolyzed water. At high electrolyzed water concentration (100%), a shrunken and rounder of the cells were conspicuously observed. Most of the cells showed signs of detachment from the surface of the well plates indicating cell death.

Antioxidant activity of electrolyzed water

The previous study has found that ERW exhibits reactive oxygen species (ROS)-scavenging activity. It can scavenge not only H₂O₂ and superoxide radical but also singlet oxygen and hydroxyl radical. ERW has an alkaline pH. In the current study, we actually observed that electrolyzed water can scavenge free radical species not only at alkaline but also at pH 2.5 and 6.0. The DPPH radical scavenging activity results are presented in Table 3 as compared with known antioxidant ascorbic acid.

According to observed experimental data in term of 50% efficient concentration (EC₅₀), unfortunately, we cannot categorize an antioxidant capacity of electrolyzed water regarding Blois' category [22]. A DPPH method could not detect radical-scavenging activity in electrolyzed water separately [23]. The intracellular ROS scavenging activity method (fluorescent stain method and immuno-spin trapping assay) and electrochemical techniques seem to be able to observe the activity specifically [23,24].

CONCLUSION

Some Indonesian people have a positive opinion about electrolyzed water marketed in Indonesia. The water has various pH with different




Fig. 2: In vitro cytotoxic activity of electrolyzed water against T47D cell lines




Fig. 3: Phase contrast images of T47D cells after incubation with 100% of electrolyzed waters at pH 2.5, 6.0, and 9 for 24 h (×20)

uses. The results provide information that electrolyzed water has antibacterial, cytotoxic, and antioxidant activities. The activity is dependent on pH. The electrolyzed water at pH 2.5 showed the highest antibacterial activity. The alkaline electrolyzed water showed the highest cytotoxicity on T47D cell lines, with an IC₅₀ of 46.6%. Furthermore, the electrolyzed water showed free radical scavenging activity with an EC₅₀ of 167.6% at pH 9.5.

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

ACKNOWLEDGMENT

The authors are highly grateful to financial support from LPPM Universitas Muhammadiyah Purwokerto.

REFERENCES

- Ignatov I, Gluhchev G, Karadzov S, Miloshev G, Ivanov N, Mosin O. Preparation of electrochemically activated water solutions (catholyte/ anolyte) and studying their physical-chemical properties. J Health Med Nurs 2015;13:64-78.
- Rahman SM, Khan I, Oh DH. Electrolyzed water as a novel sanitizer in the food industry: Current trends and future perspectives. Compr Rev Food Sci Food Saf 2016;15:471-90.
- Sharma RR, Demirci A. Treatment of *Escherichia coli* O157:H7 inoculated alfalfa seeds and sprouts with electrolyzed oxidizing water. Int J Food Microbiol 2003;86:231-7.
- Bari ML, Sabina Y, Isobe S, Uemura T, Isshiki K. Effectiveness of electrolyzed acidic water in killing *Escherichia coli* O157:H7, *Salmonella enteritidis*, and *Listeria monocytogenes* on the surfaces of tomatoes. J Food Prot 2003;66:542-8.
- Al Haq MI, Sugiyama J, Isobe S. Review applications of electrolyzed

- water in agriculture and food industries. Food Sci Technol Res 2005;11:135-50.
6. Kim MJ, Kim HK. Anti-diabetic effects of electrolyzed reduced water in streptozotocin-induced and genetic diabetic mice. Life Sci 2006;79:2288-92.
 7. Ye J, Li Y, Hamasaki T, Nakamichi N, Komatsu T, Kashiwagi T, *et al.* Inhibitory effect of electrolyzed reduced water on tumor angiogenesis. J Pharm Bull 2008;31:19-26.
 8. Tsai CF, Hsu YW, Chen WK, Ho YC, Lu FJ. Enhanced induction of mitochondrial damage and apoptosis in human leukemia HL-60 cells due to electrolyzed-reduced water and glutathione. Biosci Biotechnol Biochem 2009;73:280-7.
 9. Saitoh Y, Harata Y, Mizuhashi F, Nakajima M, Miwa N. Biological safety of neutral-pH hydrogen-enriched electrolyzed water upon mutagenicity, genotoxicity and subchronic oral toxicity. Toxicol Ind Health 2010;26:203-16.
 10. Shirahata S, Kabayama S, Nakano M, Miura T, Kusumoto K, Gotoh M, *et al.* Electrolyzed-reduced water scavenges active oxygen species and protects DNA from oxidative damage. Biochem Biophys Res Commun 1997;234:269-74.
 11. Shirahata S, Hamasaki T, Teruya K. Advanced research on the health benefit of reduced water. Trends Food Sci Technol 2012;23:124-31.
 12. Moorman E, Montazeri N, Jaykus LA. Efficacy of neutral electrolyzed water for inactivation of human norovirus. Appl Environ Microbiol 2007;73:e00653-17.
 13. Brand-Williams W, Cuvelier ME, Berset C. Use of a free radical method to evaluate antioxidant activity. Lebenswiss Technol 1995;28:25-30.
 14. Kedare SB, Singh RP. Genesis and development of DPPH method of antioxidant assay. J Food Sci Technol 2011;48:412-22.
 15. Nakagawara S, Goto T, Nara M, Ozawa Y, Hotta K, Arata Y. Spectroscopic characterization and the pH dependence of bactericidal activity of the aqueous chlorine solution. Anal Sci 1998;14:1118.
 16. Venkitanarayanan KS, Ezeike GO, Hung YC, Doyle MP. Efficacy of electrolyzed oxidizing water for inactivating *Escherichia coli* O157:H7, *Salmonella enteritidis*, and *Listeria monocytogenes*. Appl Environ Microbiol 1999;66:1476-9.
 17. McPherson LL. Understanding ORP's in the disinfection process. Water Eng Manage 1993;140:29-31.
 18. Marriott NG, Gravani RB. Principles of Food Sanitation. 5th ed. New York: Springer; 2006.
 19. Korting HC, Lukacs A, Vogt N, Urban J, Ehret W, Ruckdeschel G, *et al.* Influence of the pH-value on the growth of *Staphylococcus epidermidis*, *Staphylococcus aureus* and *Propionibacterium acnes* in continuous culture. Zentralbl Hyg Umweltmed 1992;193:78-90.
 20. Sandvik EL, McLeod BR, Parker AE, Stewart PS. Direct electric current treatment under physiologic saline conditions kills *Staphylococcus epidermidis* biofilms via electrolytic generation of hypochlorous acid. PLoS One 2013;8:e55118.
 21. O'Connor PM, Jackman J, Bae I, Myers TG, Fan S, Mutoh M, *et al.* Characterization of the p53 tumor suppressor pathway in cell lines of the national cancer institute anticancer drug screen and correlations with the growth-inhibitory potency of 123 anticancer agents. Cancer 1997;57:4285-300.
 22. Blois MS. Antioxidant determinations by the use of a stable free radical. Nature 1958;181:1199-200.
 23. Hamasaki T, Harada G, Nakamichi N, Kabayama S, Teruya K, Fugetsu B, *et al.* Electrochemically reduced water exerts superior reactive oxygen species scavenging activity in HT1080 cells than the equivalent level of hydrogen-dissolved water. PLoS One 2017;12:e0171192.
 24. Pisoschi AM, Negulescu GP. Methods for total antioxidant activity determination: A review. Biochem Anal Biochem 2011;1:1000106.

ANTIBACTERIAL, IN VITRO CYTOTOXIC, AND ANTIOXIDANT ACTIVITIES OF ELECTROLYZED OXIDIZING/REDUCING WATER

ORIGINALITY REPORT

18%

SIMILARITY INDEX

PRIMARY SOURCES

1	www.tandfonline.com Internet	158 words — 4%
2	cancerres.aacrjournals.org Internet	50 words — 1%
3	www.ncbi.nlm.nih.gov Internet	39 words — 1%
4	www.lifeionizers.com Internet	38 words — 1%
5	www.healingwatermachines.com Internet	34 words — 1%
6	Guojiao Wu, Zhiheng Chen, Peipei Wang, Mingyi Zhao et al. "Hydrogen inhalation protects hypoxic–ischemic brain damage by attenuating inflammation and apoptosis in neonatal rats", <i>Experimental Biology and Medicine</i> , 2019 Crossref	33 words — 1%
7	www.rezaghodssi.com Internet	31 words — 1%
8	greenpharmacy.info Internet	26 words — 1%
9	ffhdj.com Internet	22 words — 1%

10	www.ial.sp.gov.br Internet	18 words — 1%
11	Issa-Zacharia, A.. "In vitro inactivation of Escherichia coli, Staphylococcus aureus and Salmonella spp. using slightly acidic electrolyzed water", Journal of Bioscience and Bioengineering, 201009 Crossref	16 words — < 1%
12	www.im.microbios.org Internet	16 words — < 1%
13	www.wjgnet.com Internet	15 words — < 1%
14	Zhengxin Liu, Yongping Zheng, Jie Dang, Jiabin Zhang, Feihong Dong, Kaile Wang, Jue Zhang. "A Novel Antifungal Plasma-Activated Hydrogel", ACS Applied Materials & Interfaces, 2019 Crossref	14 words — < 1%
15	pr.hec.gov.pk Internet	14 words — < 1%
16	www.science.gov Internet	14 words — < 1%
17	ira.le.ac.uk Internet	12 words — < 1%
18	Ganyu Gu, Zhiyao Luo, Juan M. Cevallos-Cevallos, Paige Adams, George Vellidis, Anita Wright, Ariena H.C. van Bruggen. " Occurrence and population density of in irrigation ponds on produce farms in the Suwannee River Watershed ", Canadian Journal of Microbiology, 2013 Crossref	12 words — < 1%
19	refubium.fu-berlin.de Internet	12 words — < 1%
20	Se-Ri Kim, Kyoung-Yul Ryu, Myoung-Hee Lee, Chan-Sik Jung et al. "Evaluation of the Bactericidal	11 words — < 1%

Activity of Electrolyzed Water against Salmonella Typhimurium and Staphylococcus aureus on Perilla Leaves", Korean Journal of Food Science and Technology, 2012

Crossref

21 healthdocbox.com 11 words — < 1%
Internet

22 link.springer.com 9 words — < 1%
Internet

23 Castro, Inar A., Marcelo M. Rogero, Roberto M. Junqueira, and Mariana M. Carrapeiro. "Free radical scavenger and antioxidant capacity correlation of α -tocopherol and Trolox measured by three in vitro methodologies", International Journal of Food Sciences and Nutrition, 2006. 9 words — < 1%
Crossref

24 f1000.com 8 words — < 1%
Internet

25 Anil Kumar, Sanjula Baboota, SP Agarwal, Javed Ali, Alka Ahuja. "Treatment of acne with special emphasis on herbal remedies", Expert Review of Dermatology, 2014 8 words — < 1%
Crossref

EXCLUDE QUOTES OFF
EXCLUDE BIBLIOGRAPHY OFF

EXCLUDE MATCHES OFF