

# Free Radical Scavenging Activity of Juice and Ethanol Extracts of *Garcinia* *Mangostana* L. Leaves

*By* Diniatik Diniatik

2

# Free Radical Scavenging Activity of Juice and Ethanol Extracts of *Garcinia Mangostana* L. Leaves

Diniatik, Elza Sundhani, Nervita

Faculty of Pharmacy, Universitas Muhammadiyah Purwokerto

## Abstract.

The present research was conducted to assess the free radical scavenging activity of *Garcinia mangostana* L leaves by DPPH (1,1, diphenyl-2 picryl hydrazyl) method. Free radical scavenging activity of juice and ethanolic extract of *G. mangostana* leaves were determined by using spectrophotometry UV-Vis method. Juice and 96% ethanol's extract was concentrated by using rotary evaporator to get thick extract produces rendement about 2.571 % and 4.842 IC<sub>50</sub> values for juice and ethanol extract was 19.372 µg/ml and 1.965 µg/ml, it means at that concentration inhibited 50 % of free radical DPPH scavenge. IC<sub>50</sub> of d-α-tocopherol was 0.737 µg/ml. The result showed that juice are amounted ten times less than ethanol extract. When compared with d-α-tocopherol as positive control, free radical scavenging activity of juice and ethanolic extract were twenty six and three times less than d-α-tocopherol. Based on the result of anova test with significant degree 95% can be concluded that free radical scavenging activity between juice and ethanol extract were different with d-α-tocopherol.

Keywords: *Garcinia mangostana* L, free radical, DPPH, d-α-tocopherol, juice.

1

## INTRODUCTION

Free radicals are produced continually in various metabolic processes and exist in biological systems. They are important for maintaining normal physiological function. There were many report of biological activity of *G. mangostana*. From this research can conclude that free radical scavenging activity from leaves and skin bark extract of *G. mangostana* is not significantly different, if we get substance from skin bark of *G. mangostana* can damage this plant but if we get from leaf is doesn't. The aim from this research is to examined free radical scavenging activity of juice and ethanolic extract of *G. mangostana* leaves with d-α-tocopherol as standards.

## METHOD

**Plant materials.** The plants were collected from Sokaraja District, Banyumas Regency, Central Java Province, Indonesia on March 2012 and identified in laboratory of Botany and genetic Faculty of FKIP, Muhammadiyah University of Purwokerto, Indonesia. The voucher specimen was kept in Laboratory of Pharmaceutical Biology, Muhammadiyah University of Purwokerto, Indonesia. **Preparation of juice.** Leaves of *G. mangostana* (3 kg) is added with 3 liters of water at blender, after the blend was pressed with flannel cloth and was concentrated by using rotary evaporator and waterbath during five days with temperature less than 40 ° C to get thick extract produces.

**Preparation ethanol extract.** *G. mangostana* leaves were collected, dried and pulverized by using mechanical grinder. 500 grams of powder were extracted by maseration method with solvent water:ethanol 96% (1:5) during 24 hours and maseration with solvent water:ethanol 96% (1:4). After extracting the exhaustive extraction, ethanol extract was collected and concentrated under reduced pressure at 40°C by using rotary evaporator.

*Free radical scavenging activity evaluation.* Juice with concentration 8; 4; 2; and 1 µg/ml, were evaluated for its free radical scavenging activity using DPPH (diphenylpicrylhydrazyl) method, as well as Ethanol extract in serial concentration of 1; 2; 4; and 8 µg/ml. Ability to transfer electron from juice and ethanol extract were measured by its transformation from purple color to yellow color of DPPH solution in methanol observed by using UV-Vis Spectrophotometer.

There were 50 µl of juice and ethanol extract diluted on methanol and was added 0.002% DPPH solution in methanol 10 ml. After incubated at room temperature for 30 minutes, absorbance was measured at 515.6 nm using a blank solution. Free radical inhibition of DPPH in percent (%) was calculated using this formula:

$$I(\%) = \frac{Abs_{blank} - Abs_{sample}}{Abs_{blank}} \times 100$$

Where Abs<sub>blank</sub> (absorbance of blank) was absorbance of control reaction (contain all reagents without sample). Concentration where juice and ethanolic extract showing 50% inhibition (IC<sub>50</sub>) were calculated from linear regression equation from curve of percent inhibition vs. sample concentration. Antioxidant synthetics, d-α-tocopherol were used as positive control.

## RESULT

This research was designed to assess the free radicals scavenging activity from juice and ethanol extract of mangosteen leaves compared with d-α-tocopherol as standard. Extract of juice from mangosteen leaves were extracted by blending fresh leaves of mangosteen then pressing the blend with flannel cloth. The filtrate was concentrated by using rotary evaporator to get thick extract. Ethanol extract were obtained from fresh leaves of mangosteen. The leaves was washed with running water to remove any dirt or contaminants such as soil, or other materials on the leaves. Selected leaves (2 kg) were good for the next steps being aerated and dried in the sun covered with black cloth. During heating, the material is not organized piles and turned upside - behind that uniform heating and rapid drying. Drying is carried out until the moisture content is less than 10% or until the leaves and bark is easy to be destroyed when crushed. The purpose of drying is to prevent the growth of fungi or microorganisms and the decomposition of active compounds by enzymatic reaction and the hydrolysis process because of its high water content, so that crude produced is not easily damaged so it can be kept in a relatively long time. Obtained further dried simplicia was blended with blender to minimize the surface area so that the contact surface of the particles with solvent bigger bulbs and extraction becomes more optimal.


The method implemented is maceration. This method is the simplest method because it is easy to do, inexpensive, does not require sophisticated equipment. Maceration is done by soaking the powder in the liquid botanicals solvent. In this research, to improve the effectiveness of the extraction, stirring and re-maceration is conducted, macerated for 24 hours with a comparison between simplicia with 96% ethanol is 1:5, and 1:4 for re-maceration. The trick is as much as 500 grams of powder botanicals macerated with ethanol 96% as much as 2.5 liters, then pressed with flannel cloth. Obtained pulp was macerated again with 96% ethanol as much as 1.25 liters.

Extracts derived either from the juice and extract was evaporated with a rotary evaporator and over a water bath until having thick consistency. Evaporation is conducted to eliminate solvent solution so as not to affect the free radical scavenging activity assay following condensed extract obtained as shown in Table 1.

Table 1: Rendement juice and extract Mangosteen leaves

Fresh Leaves (g)	Juice		Ethanol extract		
	Thick extract (g)	Rendement (% w/w)	Simplisia (g)	thick extract (g)	Rendement (% w/w)
3000	77.14	2.571	500	96.85	4.842
(from 2 kg leaves)					

Tests conducted on the free radical scavenging activity from juice and ethanol extracts of *G. mangostana* leaves and d- $\alpha$ -tocopherol as positive control. Free radical scavenging activity was determined using DPPH (1,1-diphenyl-2-picryl hydrazil), a stable synthetic radical in aqueous solution or methanol and able to accept an electron or hydrogen radical to become a stable diamagnetic molecule. DPPH was captured by the test antioxidant that releases hydrogen, thus forming the reduced DPPH [11]. DPPH violet color changes to yellow followed by a decrease in the wavelength of maximum absorbance (515.6 nm), it showed antioxidant activity which can be seen from the% reduction.



Polyphenols that found of the *G. mangostana* L. leaves has a major role in prevention of various diseases. Polyphenols are a hundred times more effective than vitamin C and 25 times more effective than vitamin E. This substance also serves to neutralize free radicals. Leaves of *G. mangostana* contains xanthon (*mangostin*, *garsinon*) flavonoids and tannins that serves as a precursor catch (scavenger) compounds of oxygen radicals (ROS). Flavonoids have antioxidant activity that serves as a binding element in the body of harmful metals. In some plant, phenolic compounds are potent antioxidants: flavonoids, tannins, and lignins are precursors of compounds acting on the arrest of ROS.

Mechanism of action from free radical scavenging is to reduce the free radicals by giving one or more electrons to free radicals to form a normal molecule. The ability of flavonoids as antioxidants are due to the catechol group on the B ring that acts as an electron donor. Free radical scavenging activity of juice, ethanol extract (EE) and d- $\alpha$ -tocopherol were shown in Table 2. It showed that there were positive correlation between concentration of samples and percentage of inhibition. IC<sub>50</sub> value of juice was 19.372 µg/ml which was ten times less than ethanol extract (IC<sub>50</sub> 1.965 µg/ml) and twenty six times less than d- $\alpha$ -tocopherol (IC<sub>50</sub> 0.737 µg/ml). Ethanol extract was three times less

than d- $\alpha$ -tocopherol (figure 1). However, IC<sub>50</sub> of juice and ethanol extract exhibited greater value if it was compared with the IC<sub>50</sub> of ethyl acetate and acetone extract of *G. mangostana* that was reported by Sulaiman et. al. [12] (IC<sub>50</sub> 30.01 and 33.32  $\mu$ g/ml).

Table 2: IC<sub>50</sub> Value of Juice, Ethanolic extract, and d- $\alpha$ -tocopherol

Sample	Concentration (ppm)	Percentage of inhibition (%)	Linier regression equation	IC <sub>50</sub> ( $\mu$ g/ml)
Juice	100	30.906	Y = 19.	
	200	36.730	0.958. x + 37	
	400	34.466	31.560	
	800	38.511		
Ethanolic extract	100	414.239	y = 1.9	
	200	538.834	5.74.x + 7	
	400	601.982	38.4136	
	800	843.042		
d- $\alpha$ -tocopherol	20	22.006	y = 0.7	
	40	29.288	58.356x + 4	
	60	36.569	7.0456	
	80	54.854		
	100	67.579		
	19.373	0.737		

Figure 1. IC<sub>50</sub> Value of Juice, Ethanolic extract of *G. mangostana* leaves, and d- $\alpha$ -tocopherol

Juice and ethanol extract contains various compounds from *G. mangostana* L leaves parts that might have synergistic effect to scavenge free radicals from DPPH. The major secondary metabolites of mangosteen have been found to be prenylated flavonoid xanthone derivatives [13]. Prenylated flavonoid xanthone which isolated from leaves of *G. mangostana* were 1,5,8-Trihydroxy-3-methoxy-2-(3-methylbut-2-enyl)xanthone and 1,6-Dihydroxy-3-methoxy-2-(3-methyl-2-buthenyl)-xanthone (Figure 1)(5).

Based on the result of anova test with significantly degree 95%, it can be concluded that free radical scavenging activity between juice and ethanol extract were different with d- $\alpha$ -tocopherol with DPPH (1,1-difenil-2-pikril hidrazil) as free radicals.

#### CONCLUSION

Free radical scavenging activity from juice is ten times less than ethanolic extract from *G. mangostana* leaves with IC<sub>50</sub> values for juice and ethanolic extract was 19.372 and 1.965  $\mu$ g/ml with DPPH as free radicals and it needs further confirmation of compounds that is contained in juice and ethanolic extract.

#### ACKNOWLEDGEMENT

The researchers would like to thank Directorate General of Higher Education, Ministry of National Education Republic of Indonesia for the research funding.

# Free Radical Scavenging Activity of Juice and Ethanol Extracts of Garcinia Mangostana L. Leaves

ORIGINALITY REPORT

# 10%

SIMILARITY INDEX

## PRIMARY SOURCES

- 1** [www.japer.in](http://www.japer.in)  
Internet 36 words — 2%
- 2** Manosroi, J., R. Wilairat, A. Kijjoa, and A. Manosroi. "Free Radical Scavenging Activity of Extracts from Thai Plants in Guttiferae and Schisandraceae Families", *Pharmaceutical Biology*, 2005.  
Crossref 34 words — 2%
- 3** "Garcinia binucao crude ethanolic leaf extract prevents alcohol-induced neurotoxic effects on learning, short-term memory, and motor functions in *Drosophila melanogaster*", *Journal of Applied Pharmaceutical Science*, 2018  
Crossref 26 words — 1%
- 4** I Made Ginarsa, Adi Soeprijanto, Mauridhi Hery Purnomo, Syafaruddin, Takashi Hiyama. "Controlling voltage collapse using ANFIS-based composite controller-SVC in power systems", *TENCON 2011 - 2011 IEEE Region 10 Conference*, 2011  
Crossref 17 words — 1%
- 5** [japanesegreenteaonline.com](http://japanesegreenteaonline.com)  
Internet 14 words — 1%
- 6** [pharmacologyonline.silae.it](http://pharmacologyonline.silae.it)  
Internet 14 words — 1%
- 7** Dundar, A., O. F. Yesil, H. Acay, V. Okumus, S. Ozdemir, and A. Yildiz. "Antioxidant properties,

chemical composition and nutritional value of *Terfezia boudieri* (Chatin) from Turkey", Food Science and Technology International, 2012.

Crossref

8

[www.omicsonline.org](http://www.omicsonline.org)

Internet

12 words — 1%

9

[www.mdpi.com](http://www.mdpi.com)

Internet

9 words — < 1%

10

Dmitriy Obolskiy. "*Garcinia mangostana* L.: a phytochemical and pharmacological review", Phytotherapy Research, 08/2009

Crossref

7 words — < 1%

EXCLUDE QUOTES OFF

EXCLUDE MATCHES OFF

EXCLUDE BIBLIOGRAPHY OFF